

3D-HST+CANDELS: THE EVOLUTION OF THE GALAXY SIZE-MASS DISTRIBUTION SINCE $Z = 3$

A. VAN DER WEL¹, M. FRANX², P.G. VAN DOKKUM³, R.E. SKELTON⁴, I.G. MOMCHEVA³, K.E. WHITAKER⁵,
G.B. BRAMMER⁶, E.F. BELL⁷, H.-W. RIX¹, S. WUYTS⁸, H.C. FERGUSON⁶, B.P. HOLDEN⁹, G. BARRO⁹,
A.M. KOEKEMOER⁶, YU-YEN CHANG¹, E.J. MCGRATH¹⁰, B. HÄUSSLER^{11 12}, A. DEKEL¹³, P. BEHROOZI⁶, M. FUMAGALLI²,
J. LEJA³, B.F. LUNDGREN¹⁴, M.V. MASEDA¹, E.J. NELSON³, D.A. WAKE^{14 15}, S.G. PATEL¹⁶, I. LABBÉ², S.M. FABER⁹,
N.A. GROGIN⁶, AND D.D. KOCEVSKI¹⁷

Accepted for publication in ApJ

ABSTRACT

Spectroscopic+photometric redshifts, stellar mass estimates, and rest-frame colors from the 3D-HST survey are combined with structural parameter measurements from CANDELS imaging to determine the galaxy size-mass distribution over the redshift range $0 < z < 3$. Separating early- and late-type galaxies on the basis of star-formation activity, we confirm that early-type galaxies are on average smaller than late-type galaxies at all redshifts, and find a significantly different rate of average size evolution at fixed galaxy mass, with fast evolution for the early-type population, $R_{\text{eff}} \propto (1+z)^{-1.48}$, and moderate evolution for the late-type population, $R_{\text{eff}} \propto (1+z)^{-0.75}$. The large sample size and dynamic range in both galaxy mass and redshift, in combination with the high fidelity of our measurements due to the extensive use of spectroscopic data, not only fortify previous results, but also enable us to probe beyond simple average galaxy size measurements. At all redshifts the slope of the size-mass relation is shallow, $R_{\text{eff}} \propto M_*^{0.22}$, for late-type galaxies with stellar mass $> 3 \times 10^9 M_\odot$, and steep, $R_{\text{eff}} \propto M_*^{0.75}$, for early-type galaxies with stellar mass $> 2 \times 10^{10} M_\odot$. The intrinsic scatter is $\lesssim 0.2$ dex for all galaxy types and redshifts. For late-type galaxies, the logarithmic size distribution is not symmetric, but skewed toward small sizes: at all redshifts and masses a tail of small late-type galaxies exists that overlaps in size with the early-type galaxy population. The number density of massive ($\sim 10^{11} M_\odot$), compact ($R_{\text{eff}} < 2\text{kpc}$) early-type galaxies increases from $z = 3$ to $z = 1.5 - 2$ and then strongly decreases at later cosmic times.

1. INTRODUCTION

The size distribution of the stellar bodies of galaxies, and its evolution with cosmic time, provides important

clues about the assembly history of galaxies and the relationship with their dark matter halos. The two main classes of galaxies, early- and late types, show very different dependencies between size and stellar mass (Shen et al. 2003). The weak dependence between size and mass for late-type galaxies implies that the high-mass late types, on average, have higher surface mass densities than low-mass late types. In contrast, early types show a more complex relationship between stellar mass and density, with the density peaking for systems with stellar masses around $M_* \sim 4 \times 10^{10} M_\odot$ and decreasing towards both lower and higher masses, as reflected in the classical Kormendy (1977) relation. This fundamental difference does not depend on whether classification of early and late types is based on star formation activity, bulge dominance (Sérsic index) or visual inspection, and implies that the two types have very different evolutionary and assembly histories.

In this paper we present the evolution of the size-mass distribution up to $z = 3$, based on spectroscopy and multi-wavelength photometry from the 3D-HST survey (Brammer et al. 2012a) and HST/WFC3 imaging from CANDELS (Grogin et al. 2011; Koekemoer et al. 2011). Angular galaxy sizes are measured from the CANDELS imaging as described by van der Wel et al. (2012) and the HST/WFC3 grism observations from 3D-HST provide spectroscopic confirmation and redshifts for a large fraction of the sample, considerably strengthening, with respect to previous studies, the fidelity of estimates for stellar masses and rest-frame photometric properties.

So far, most previous studies have focused on the evolution of average galaxy sizes of the high-mass end of

¹ Max-Planck Institut für Astronomie, Königstuhl 17, D-69117, Heidelberg, Germany; e-mail: vdwel@mpia.de

² Leiden Observatory, Leiden University, P.O.Box 9513, NL-2300 AA Leiden, Netherlands

³ Department of Astronomy, Yale University, New Haven, CT 06511, USA

⁴ South African Astronomical Observatory, PO Box 9, Observatory, 7935, South Africa

⁵ Astrophysics Science Division, Goddard Space Center, Greenbelt, MD 20771, USA

⁶ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁷ Department of Astronomy, University of Michigan, 500 Church Street, Ann Arbor, MI 48109, USA

⁸ Max-Planck Institut für Extraterrestrische Physik, Giessenbachstrasse, D-85748 Garching, Germany

⁹ University of California Observatories/Lick Observatory, University of California, Santa Cruz, CA 95064, USA

¹⁰ Department of Physics and Astronomy, Colby College, Waterville, ME 0490, USA

¹¹ Physics Department, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

¹² Centre for Astrophysics, Science & Technology Research Institute, University of Hertfordshire, Hatfield, Herts, AL10 9AB, UK

¹³ Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

¹⁴ Department of Astronomy, University of Wisconsin, Madison, WI 53706, USA

¹⁵ The Department of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK

¹⁶ Observatories of the Carnegie Institution of Washington, Pasadena, CA 91101, USA

¹⁷ Department of Physics and Astronomy, University of Kentucky, Lexington, KY 40506, USA

the distribution ($\gtrsim 5 \times 10^{10} M_{\odot}$). Enabled by both the improved data quality and a five-fold increase in sample size, we can now, for the first time, describe the size distribution of galaxies across redshift.

1.1. Size Evolution of Late-Type Galaxies

Tracing the evolution of the size distribution with redshift allows us to test the most basic elements in our theory of galaxy formation. The zero-th order expectation is that disk scale lengths evolve fast, approximately as the inverse of the Hubble parameter (Mo, Mao & White 1998), and early and recent work on the average sizes of Lyman Break Galaxies (LBGs) at high redshifts ($z \sim 2 - 6$) roughly agree with this expectation for a Λ CDM cosmology: Giavalisco et al. (1996), Ferguson et al. (2004), Oesch et al. (2010), and Mosleh et al. (2012) all find rapid size evolution with redshift: $R_{\text{eff}} \propto (1+z)^{\beta=-1.1}$.

In contrast, the average size at a given stellar mass of the population of disk-dominated galaxies evolves slowly at late times ($z \lesssim 1$) has been reported to evolve slowly as measured at fixed galaxy mass ($\beta = -0.2$) or not at all (Lilly et al. 1998; Ravindranath et al. 2004; Barden et al. 2005). The implication would be that the evolution of the disk galaxy population is decoupled from the evolution of the dark matter halo population. One fundamental difference between the results on LBGs and lower-redshift disk galaxies is the rest-frame wavelength at which the sizes are measured: the rest-frame UV light seen for LBGs originates from young stars that may be, and are generally expected to be, distributed differently than bulk of baryonic and stellar mass, not to mention the consequences of extinction.

The advent of ground-based near-infrared imaging surveys helped to bridge the $z < 1$ and $z > 2$ regimes by enabling size measurements in a consistent manner at a fixed rest-frame wavelength. Early results suggested slow evolution for late-type galaxies up to $z \sim 3$ (Trujillo et al. 2006a), but the uncertainties at $z > 1$ were such that evolution in that regime was not strongly constrained. Later ground-based work pointed at faster evolution at a fixed galaxy mass: Franx et al. (2008) found $\beta = -0.6$, and Williams et al. (2010) found $\beta = -0.9$, but precise constraints at $z > 1.5$ remained elusive and the apparent tension between the $z \lesssim 1$ work and the near-infrared at $z \sim 1.5$ unaddressed.

Several HST/NICMOS-based studies of the morphology and structure of massive $z \sim 2$ galaxies in the rest-frame optical eventually led to mostly converged results, with $\beta \sim -0.8$ for massive ($\sim 10^{11} M_{\odot}$), star-forming galaxies from $z \sim 2.5$ to the present (Toft et al. 2007; Buitrago et al. 2008; Kriek et al. 2009b). Whether the difference with the significantly faster evolution for LBG galaxies ($\beta \sim -1.1$) is caused by morphological K corrections, the difference in mass (the typical LBG has $M_{*} \sim 10^{10} M_{\odot}$), or physical changes with redshift has so far remained unclear. In addition, the difference with the previously mentioned studies at $z < 1$ (Lilly et al. 1998; Ravindranath et al. 2004; Barden et al. 2005) remains unexplained.

Improving the measurement of β and its mass-dependence is crucial to make the next step in understanding disk galaxy formation. In this paper we will

address these issues and describe the full size-mass distribution of high-redshift galaxies over a broad range in galaxy mass and redshift. We will

- measure the evolution of the slope of the size-mass relation;
- present the size distribution as a function of stellar mass and redshift;
- provide a consistent comparison with UV-selected, high-redshift samples.

1.2. Size Evolution of Early-Type Galaxies

Over the past five years, more attention has been bestowed on the size evolution of early-type galaxies than on the size evolution of late-type, star-forming galaxies. Interest in the topic was initiated by reports that $z \sim 1.5$ early-type galaxies have remarkably small sizes in HST-based, rest-frame UV imaging (Daddi et al. 2005; Trujillo et al. 2007) and ground-based near-infrared imaging (Trujillo et al. 2006b). NICMOS imaging presented by Zirm (2007), Toft et al. (2007), Stockton et al. (2008), and McGrath et al. (2008) provided space-based, rest-frame optical size measurements which strengthened the evidence for rapid size evolution ($\beta = -1$ or faster) as measured at a fixed galaxy mass ($\sim 10^{11} M_{\odot}$). This notion became firmly established through larger samples (Buitrago et al. 2008) and the first spectroscopic samples (van Dokkum et al. 2008; Damjanov et al. 2011).

Concerns regarding gross overestimates of the stellar mass content of the compact early-type galaxies were alleviated by dynamical mass estimates of such galaxies at $z \gtrsim 1$ (van der Wel et al. 2008; Cimatti et al. 2008; Newman et al. 2010; van de Sande et al. 2011; Toft et al. 2012; van de Sande et al. 2013; Bezanson et al. 2013; Belli et al. 2014), and the analysis by Szomoru et al. (2010) of ultra-deep imaging of a single compact galaxy has demonstrated the absence of low-surface brightness wings that could have been missed by shallower imaging. Increases in sample size and dynamic range in stellar mass have constrained the average size evolution of early-type galaxies with stellar masses $> 5 \times 10^{10} M_{\odot}$ to $\beta \sim -1.3$ up to $z = 2.5$, with no evidence for a change in the slope of the relation over this mass range (Newman et al. 2012). The steepness of the relation combined with the non-negligible scatter accommodates observations that early-type galaxies display a large range in size at $z > 1$ (e.g., Mancini et al. 2010; Saracco et al. 2011).

While the observational results have largely converged, the interpretation is still debated. Some authors have considered the average increase in size over time due to the addition of new, larger early-type galaxies, and while some argue that this cannot reproduce the observations (van der Wel et al. 2009a), others argue that a population of compact early-type galaxies (with sizes $R_{\text{eff}} \lesssim 2\text{kpc}$) exists within present-day clusters, with a number density comparable to that of higher-redshift early-type galaxies (Valentinuzzi et al. 2010; Poggianti et al. 2013); tension with the absence of such galaxies in the Sloan Digital Sky Survey (SDSS) remains (Trujillo et al. 2009; Taylor et al. 2010). The crucial observational test is to trace the evolution of the number density of early-type galaxies as a function of their size. Cassata et al. (2011),

Szomoru et al. (2012), and Newman et al. (2012) show strong evolution in the number density of small galaxies at $0 < z < 2.5$, while Carollo et al. (2013) claim no evolution at $0 < z < 1$. Our use of 5 fields address the issue of field-to-field variations that may affect the aforementioned studies based on smaller samples, and extend the dynamic redshift range of the Carollo et al. sample.

The leading explanation for the size growth of individual galaxies is accretion and tidal disruption of satellite galaxies that gradually build up the outer parts. For this process, the change in size is large compared to the increase in mass: $\Delta R_{\text{eff}} \propto \Delta M_*^2$ (e.g., Bezanson et al. 2009; Hopkins et al. 2009). This analytical prediction based on conservation of binding energy has been tested through numerical simulations (e.g., Naab et al. 2009; Oser et al. 2012; Bédorf & Portegies Zwart 2013). The analytically predicted and simulated evolution in the increased surface mass density at large radii is, in fact, observed (van Dokkum et al. 2010); in addition, the central stellar density shows little evolution (Bezanson et al. 2009; Hopkins et al. 2009; van Dokkum et al. 2010), also consistent with a minor-merger scenario. In other words, the observations show that there is no need and little room to physically expand a galaxy by displacing large numbers of stars to large radii through rapid changes in the central potential, as suggested by Fan et al. (2010). A possible challenge to the minor merger scenario is posed by the lack of strong evolution in the slope of the mass density profile seen in lensing galaxies (Sonnenfeld et al. 2013).

Until recently, the size evolution of late- and early-type galaxies was usually discussed separately and treated as different topics. However, in order to understand the joint evolution of these classes one has to take into account the continuous transition of late-type to early-type galaxies seen in particular in the stellar mass range of $10^{10} M_\odot$ to $10^{11} M_\odot$ (e.g., Bell et al. 2004; Faber et al. 2007; Brown et al. 2007; Ilbert et al. 2010; Brammer et al. 2011; Buitrago et al. 2013; Muzzin et al. 2013). The star-forming progenitors of the small early-type galaxies are now looked for and plausibly identified (e.g., Whitaker et al. 2011; Barro et al. 2013,?; Toft et al. 2014), but the evolutionary path of the transitioning galaxies has not been fully mapped out.

In this paper, regarding the evolution of early-type galaxies, we will

- present the distribution of sizes as a function of stellar mass and redshift, jointly with those of late-type galaxies;
- show the evolution of the number density of early-type galaxies as a function of size.

After describing the data and sample selection in §2 we present and analyze size distributions as a function of redshift and galaxy mass in §3. We compare our results with previous studies in §4 and then discuss the implications of our findings in §5. We assume the cosmological parameters $(\Omega_M, \Omega_\Lambda, h) = (0.27, 0.73, 0.71)$. Finally, we use AB magnitudes and the Chabrier (2003) stellar initial mass function.

2. DATA

The procedures for source detection, multi-wavelength photometry, redshift determinations, rest-frame color and stellar mass estimates are described elsewhere. Here, we briefly summarize these steps.

2.1. Source Detection

207,967 sources in all five CANDELS/3D-HST fields (Koekemoer et al. 2011; Brammer et al. 2012a) are detected in and extracted from images that combine the available HST/WFC3 IR channel data; that is, stacked mosaics consisting of F125W, F140W and F160W imaging are constructed for this purpose. We refer to the photometry data release paper from the 3D-HST collaboration by Skelton et al. (2014) for details.

2.2. Photometric and Spectroscopic Redshift Determinations

Multi-wavelength photometry from HST/WFC3 and HST/ACS imaging is produced by creating PSF-matched images, using custom-made kernels and performing simple aperture photometry. Multi-wavelength photometry from ground-based optical-NIR and Spitzer/IRAC imaging is produced using the approach outlined by Labbé et al. (2006) and further developed by González et al. (2010), which addresses blending by nearby sources and takes the large differences in PSF into account through the use of custom-made convolution kernels. Like alternative methods such as TFIT (Laidler et al. 2007), our approach uses high-resolution images as priors to model sources in lower-resolution images.

Photometric redshifts are determined based on the multi-wavelength photometry using the EAZY package (Brammer et al. 2008). (Skelton et al. 2014) describe the procedure in full, but it essentially follows Whitaker et al. (2011): briefly, linear combinations of a set of templates that span the range of observed galaxy properties are used to fit the photometry, producing a marginalized, posterior probability distribution for the redshift, with z_{phot} as its peak.

The photometric redshifts provide a baseline for 3D-HST WFC3 G141 grism spectroscopy to provide more precise redshift information. Brammer et al. (2012a) describe the extraction of spectra and redshift determination in detail. The method has been updated to use interlaced rather than drizzled HST/WFC3 mosaics, which are used to construct the photometric catalogs. For all sources brighter than $F_{160W, \text{AB}} = 23$, the F140W image is traced along the dispersed WFC3/G141 grism image, such that a spectrum is extracted that accounts for the convolved spectral and spatial information of the low wavelength resolution ($R \sim 130$) grism data. For each extracted object the spectroscopic information is combined with the photometric redshift probability distribution, producing a new, best-fitting redshift (Momcheva et al. in prep.). Finally, spectroscopic redshifts from the literature are used when available.

The grism data significantly improve the redshift precision for thousands of galaxies, and provide indispensable evidence for the good accuracy of the purely photometric redshift estimates. The current version of the 3D-HST redshift catalog contains grism redshift information for all objects brighter than $H_{F160W} = 23$ and for which such data is available ($\sim 75\%$ of the CANDELS




Figure 1. Rest-frame $U - V$ vs. $V - J$ color distribution for six redshift bins (each 0.5 wide). The two distinct classes of quiescent and star-forming galaxies are separated by the indicated selection criteria to define our early- and late-type galaxy samples.

area). For our mass-limited sample – defined below in Section 2.4 – this amounts to $\sim 10,000$ galaxies. This is $\sim 30\%$ of the total sample and 50% for the sample of massive ($M_* > 10^{10} M_\odot$) galaxies in the crucial redshift range $1 < z < 3$. For these galaxies the Quadri & Williams (2010) pair test demonstrates a precision of $\Delta z / (1 + z) = 0.003$, or 0.3% . For purely photometric redshifts this is $1 - 2.5\%$, depending on the varying photometric dataset available for each of the five fields, suggesting a factor $3 - 10$ improvement in redshift precision from the grism data. There is no systematic offset between the two sets of redshifts.

2.3. Rest-Frame Colors and Stellar Mass Estimates

EAZY is used to compute the rest-frame $U - V$ and $V - J$ colors and the package FAST (Kriek et al. 2009a) is used to estimate stellar masses. A large number of Bruzual & Charlot (2003) templates with solar metallicity, a wide range in age (4×10^7 yr to 12.5×10^9 yr, but always younger than the universe), exponentially declining star-formation histories (with time scales $\tau = 10^7 - 10^{10}$ yr), and dust extinction ($A_V = 0 - 4$) are used and matched to the photometry. The final stellar mass is corrected for the difference between the total F160W flux from the photometric catalog and the total F160W as measured with GALFIT (see §2.5). F125W is used in case F160W is not available. This correction ensures that our size and mass estimates are mutually consistent in that both are based on the same model for the light distribution.

The uncertainties in the stellar mass estimates can be large and are to some extent still unknown. However,

the possible error in mass-to-light ratio for low-mass blue galaxies with precisely known redshifts due to uncertainties in the SFH and metallicity is not larger than a factor of a few. Moreover, for high-mass galaxies the relation between dynamical and stellar mass estimates is the same to within a factor ~ 2 over the whole redshift range $0 < z < 2$, indicating that stellar masses are estimated reasonably consistently at different redshifts (e.g., van der Wel et al. 2006; van de Sande et al. 2013). The direct mass measurement of a $z \sim 1.5$ strong gravitational lens is also in agreement with the photometrically estimated stellar mass (van der Wel et al. 2013). For these reasons we are confident that the analysis presented in this paper, which looks at galaxies that span several orders of magnitude in stellar mass, is not fundamentally dependent on the current uncertainties in the mass estimates.

2.4. Sample Selection

Following Wuyts et al. (2007) and Williams et al. (2009), we utilize rest-frame colors to distinguish between the two basic classes of galaxies: star-forming and quiescent galaxies. In this paper we refer to the former as late-type galaxies, and to the latter as early-type galaxies. In Figure 1 we show the rest-frame $U - V$ and $V - J$ color distribution of galaxies with stellar mass in excess of $10^{10} M_\odot$ for a range of redshifts. Even beyond $z = 2$ the early- and late-type galaxies are separated in this space, which allows us to effectively assign each object to one of the two classes.

An indication of the high fidelity of this selection method is that less than 20% of the thus classified early-




Figure 2. Observed H_{F160W} magnitude versus stellar mass in six redshift bins. The color-coding represents the rest-frame $U - V$ colors, ranging from $U - V = 0$ (blue) to $U - V = 2$ (red). The horizontal dashed lines indicate the limit ($H_{F160W} = 24.5$) down to which we can determine sizes with good fidelity. The vertical lines illustrate the resulting mass completeness limits for blue ($U - V = 0.5$) and red ($U - V = 2.0$) galaxies, respectively. See §2.4 for further details.

type galaxies with matches in the FIREWORKS catalog from Wuyts et al. (2007) are detected at $24\mu\text{m}$, while more than $\sim 80\%$ of the late-type galaxies are detected – these numbers are for the mass range $3 \times 10^{10} M_\odot < M_\odot < 10^{11} M_\odot$ and the redshift range $1 < z < 2$. We note that a simpler selection by just $U - V$ color would compromise the separation into types, as the sub-sample of galaxies that are red in $U - V$ consist of dusty and quiescent objects. In addition, 80% of the color-selected early-type (late-type) galaxies have Sérsic indices large (smaller) than $n = 2.5$.

We adopt redshift- and color-dependent mass limits (Figure 2) that are set by the $F160W$ magnitude limit down to which galaxy sizes – here described in §2.5 – can be determined with high fidelity. van der Wel et al. (2012) showed that galaxy sizes can be precisely and accurately determined down to a magnitude limit of $H_{F160W} = 24.5$; at fainter magnitudes the random and systematic errors can exceed 20% for large galaxies with large Sérsic indices. Since most $z > 1$ galaxies in our sample have small sizes ($0.3''$ in the median) and low Sérsic indices (1.4 in the median) this magnitude limit is conservative.

Out to $z = 3$ our magnitude-limited sample is complete in stellar mass down to $\sim 10^{10} M_\odot$ (see Figure 2). This limit is 1.9 magnitudes brighter than the 5σ detection limit, and simulations of artificial objects inserted in the images show that $\gtrsim 95\%$ of all objects are detected (Skelton et al. 2014). Therefore, incompleteness will not be an issue for our sample, and biases against large (or

small) galaxies will not play a significant role.

Our mass-selected sample, with a redshift-dependent mass limit as described above (and a minimum of $M_* = 10^9 M_\odot$ and $H_{F160W} = 25.5$ at all redshifts), contains 32,722 galaxies. 43 of these are excluded because of catastrophically failed surface brightness profile fits. We then manually verified the SEDs, size measurements, mass and redshift estimates of all 7,065 objects with flagged GALFIT results ($f = 2$ from van der Wel et al. 2012), small or large sizes ($R_{\text{eff}} < 0.6$ kpc; $R_{\text{eff}} > 10$ kpc), large stellar masses ($M_*/M_\odot > 10^{10.8}$), or large differences between photometric and grism redshifts ($\Delta z > 0.15$), as well as all early-type galaxies at redshifts $2 < z < 3$. 1,721 (5.3%) were removed after this verification due to errors in the size, redshift, or mass measurements. Hence, we have a final sample of 30,958 galaxies; 21,828 (5,189) are at $z > 1$ ($z > 2$).

In Figure 3 we show the rest-frame $(U - V) - M_*$ distribution – a clear bi-modality is seen, equivalent to the bi-modality seen in Figure 1. As is well-known, more massive galaxies are redder and more likely to be quiescent, at least up to $z \sim 2.5$. The most massive galaxies $> 10^{11} M_\odot$ are essentially all red. They are a mix of quiescent and star-forming galaxies; the quiescent galaxies dominate in number at $z < 1$; the star-forming galaxies at $z > 2$.

2.5. Size Determinations

Galaxy sizes are measured as described by van der Wel et al. (2012) from mosaics created by the 3D-HST col-




Figure 3. Rest-frame $U - V$ color vs. stellar mass in six redshift bins. Early-type galaxies, defined as illustrated in Figure 1, are shown in red; late types in blue. A clearly defined red sequence is seen up to $z = 3$, with an increased prevalence of dusty late-type galaxies toward higher redshifts.

laboration from public CANDELS F125W and F160W and 3D-HST F140W raw imaging data. With GALFIT we fit single-component Sérsic profiles to the 2-dimensional light profiles of all detected objects, making use of custom-made PSF models; the package GALAPAGOS allows for simultaneous fitting of as many neighboring objects as needed, in order to avoid confusion. As the effective radius we use the semi-major axis of the ellipse that contains half of the total flux of the best-fitting Sérsic model. For a full description of the procedure we refer to van der Wel et al. (2012); the size measurements published in their catalog and the size measurements here¹⁸ are fully consistent.

Color gradients and their evolution affect the size measurements to a degree that greatly exceeds the statistical uncertainties in our sample averages and the size distributions as a function of galaxy mass and redshift. The dynamic range in wavelength for our sample is rather small (from $1.25\mu\text{m}$ to $1.6\mu\text{m}$) which does not allow to systematically probe the effect of color gradients over the large range in redshift. To extend the dynamic range we analyze ACS/F814W imaging in the COSMOS field from CANDELS parallel observations. GALAPAGOS is used to obtain galaxy sizes in precisely the same manner for the F814W data as for the WFC3 data.

Using this extension in wavelength we show in Fig-

ure 4 the wavelength dependence of galaxy size for a sample of 777 late-type galaxies. The figure shows $\Delta \log R_{\text{eff}} / \Delta \log \lambda$ as a function of galaxy mass and redshift. For galaxies at $z < 1$, the difference between $\log R_{\text{eff}}$ from the F814W and F125W size measurements was taken as $\Delta \log R_{\text{eff}}$; for $z > 1$ galaxies the difference between $\log R_{\text{eff}}$ from the F125W and F160W measurements were taken. The pivot wavelengths of the respective filters are used to compute $\Delta \log \lambda$.

The generally negative values of the quantity $\Delta \log R_{\text{eff}} / \Delta \log \lambda$ imply that late-type galaxies are typically smaller at longer wavelengths and thus have negative color gradients (redder centers). The color gradients of $z > 1$ galaxies have been extensively studied before (e.g., Szomoru et al. 2011; Wuyts et al. 2012), and here we merely mention the trends that are relevant for our conversion of the measured R_{eff} to the R_{eff} at a rest-frame wavelength of 5000\AA . Color gradients are strongest for the most massive galaxies at all redshifts, and stronger at later cosmic times for galaxies of all masses. The evolution is remarkably smooth, which allows us to parametrize the wavelength-dependence of R_{eff} as a simple function of redshift and galaxy stellar mass:

$$\frac{\Delta \log R_{\text{eff}}}{\Delta \log \lambda} = -0.35 + 0.12z - 0.25 \log \left(\frac{M_*}{10^{10} M_{\odot}} \right). \quad (1)$$

Kelvin et al. (2012) derive a very similar wavelength dependence of R_{eff} , but a direct comparison cannot be made as those authors do not distinguish between galax-

¹⁸ The results from our GALFIT profile fits used here, with IDs matched to the Skelton et al. (2014) catalogs, are publicly available for all 5 fields:

<http://www.mpia-hd.mpg.de/homes/vdwel/3dhstcandels.html>.




Figure 4. Wavelength dependence of R_{eff} in bins of stellar mass and redshift; the latter is indicated by the color coding. Late-type galaxies, as defined in Figure 1, with robust size measurements from ACS/F814W, WFC3/F125W and WFC3/F160W imaging are included (see text for details). Generally, sizes are smaller at longer wavelengths, that is, late-type galaxies are bluer in the outer parts. Moreover, this gradient is stronger for more massive galaxies at all redshifts, and the gradient decreases with redshift, at the same rate for all masses. The dotted lines represent the parametrization given in Eq. 1 that we use to correct our size measurements of late-type galaxies.

ies with different masses. Then, R_{eff} at a rest-frame wavelength of 5000\AA is estimated as

$$R_{\text{eff}} = R_{\text{eff},F} \left(\frac{1+z}{1+z_p} \right)^{\frac{\Delta \log R_{\text{eff}}}{\Delta \log \lambda}}, \quad (2)$$

where F denotes either F125W (for galaxies at $z < 1.5$) or F160W (for galaxies at $z > 1.5$), and z_p is the ‘pivot redshift’ for these respective filters (1.5 for F125W and 2.2 for F160W). Positive color gradients as computed with Equation 1 are set at zero.

A similarly detailed correction for early-type galaxies is currently not feasible as that sub-sample is much smaller and has much redder colors. For the 122 early-type galaxies in the COSMOS field at redshifts $0 < z < 2$ and with robust size measurements in all three filters, we find an average size gradient of $\Delta \log R_{\text{eff}} / \Delta \log \lambda = -0.25$, with no discernible trends with mass and redshift, in reasonable agreement with Kelvin et al. (2012) for low-redshift and Guo et al. (2011) for high-redshift early-type galaxies. Hence, we adopt this value for all early-type galaxies when computing their rest-frame 5000\AA sizes with Equation 2.

This rather convoluted procedure has a small but significant effect on the size measurements, with implications for the rate of size evolution. In the redshift range $1 < z < 2$ the size corrections are always smaller than 0.05 dex, because the observed wavelength is always close to the desired rest-frame wavelength; at $z < 1$ the correction can increase up to 0.15 dex, but there the color gradient is well-constrained (see Figure 4) for all galaxy

masses.

3. EVOLUTION OF THE SIZE-MASS DISTRIBUTION AT $0 < z < 3$

The size-mass distributions for early- and late-type galaxies as a function of redshift are shown in Figure 5. The first basic observation is that a well-defined size-mass relation exists at all redshifts $0 < z < 3$ for both populations. With increasing redshift, the two classes have increasingly different size-mass relations, mostly as a result of strongly decreasing R_{eff} for early types. In addition, at all redshifts the slope of the relation is steeper for early types than for late types. At all stellar masses $\lesssim 10^{11} M_{\odot}$ early-type galaxies are, on average, smaller than late-type galaxies, but at very high stellar masses ($> 2 \times 10^{11} M_{\odot}$) the two populations have similar sizes at all redshifts, as far as the small samples allow for such a comparison.

However, the relation for early-type galaxies flattens out below $M_* = 2 \times 10^{10} M_{\odot}$, at least up to $z = 1.5$ beyond which our sample is incomplete at these low masses. This implies that the peak in surface mass density at $\sim 4 \times 10^{10} M_{\odot}$ seen in the present-day universe also existed at larger look-back times, at least up to $z \sim 2$.

In §3.1 we will first provide an analytical description of the size-mass relation, which allows us to take cross-contamination between the two types and outliers into account. In the remainder of this section we will provide various direct measurements such as median sizes and percentile distributions as a function of mass and redshift, and describe trends that are not captured by our analytical description, such as deviations from a single power law and skewness of the size distribution.

3.1. Analytical Description

The basic characteristics of the galaxy size distribution are given by the slope, intercept and (intrinsic) scatter of size as a function of mass. We parametrize this following Shen et al. (2003) and assume a log-normal distribution $N(\log r, \sigma_{\log r})$, where $\log r$ is the mean and $\sigma_{\log r}$ is the dispersion. Furthermore, r is taken to be a function of galaxy mass:

$$r(m_*)/\text{kpc} = A \cdot m_*^{\alpha}, \quad (3)$$

where $m_* \equiv M_*/7 \times 10^{10} M_{\odot}$. As we will describe in §3.3 it is reasonable to assume that $\sigma_{\log r}$ is independent of mass.

The model distribution $N(\log r(m_*), \sigma_{\log r})$ prescribes the probability distribution for observing R_{eff} for a galaxy with mass m_* . If the measured R_{eff} has a Gaussian, $1-\sigma$ uncertainty of $\delta \log R_{\text{eff}}$, then the probability for this observation is the inner product of two Gaussians:

$$P = \langle N(\log R_{\text{eff}}, \delta \log R_{\text{eff}}), N(\log r(m_*), \sigma_{\log r}) \rangle. \quad (4)$$

Thus we compute for each galaxy the probabilities P_{ET} and P_{LT} for the respective size-mass distribution models for the early-type and late-type populations. Incompleteness terms should formally be included in these probabilities (as described by, e.g., Huang et al. 2013), but due to our conservative sample selection (see §2.4) we are not biased against faint, large objects.

The uncertainty in size, $\delta \log R_{\text{eff}}$, is computed as outlined by (van der Wel et al. 2012). A random uncertainty




Figure 5. Size-stellar mass distribution of late- and early-type galaxies (same symbols as in Figure 2). A typical 1σ error bar for individual objects in the higher-redshift bins is shown in the bottom-right panel. The lines indicate model fits to the early- and late-type galaxies as described in §3.1. The dashed lines, which are identical in each panel, represent the model fits to the galaxies at redshifts $0 < z < 0.5$. The solid lines represent fits to the higher-redshift samples. The mass ranges used in the fits are indicated by the extent of the lines in the horizontal direction. Strong evolution in the intercept of the size-mass relation is seen for early-type galaxies; moderate evolution for the late-type galaxies (also see Figure 6). There is no significant evidence for evolution in the slope (also see Figure 6). The parameters of the fits shown here are given in Table 1.

of 0.15 dex in m_* is included in our analysis by treating it as an additional source of uncertainty in R_{eff} : for a size-mass relation with a given slope, an offset in m_* translates into an offset in R_{eff} . Hence, the calculation of P stays 1-dimensional. The fiducial slopes we use to convert $\delta \log R_{\text{eff}}$ into δm_* are $\alpha = 0.7$ for early-type galaxies and $\alpha = 0.2$ for late-type galaxies.

We also take into account the mis-classification of early- and late-type galaxies. Despite the bimodal distribution in the color-color diagram (§2.4; Figure 1) there are galaxies in the region between the star-forming and quiescent sequences, making their classification rather arbitrary, causing cross-contamination of the two classes (also see Holden et al. 2012). Motivated by this work, we take this mis-classification probability to be 10%. We will comment on the effects of varying this parameter below, when we describe the fitting results.

The mis-classification probability precisely corresponds to the early- and late-type contamination fractions in a sample in case the two sub-samples have equal numbers of galaxies. The actual contamination fraction scales with the early-/late-type fraction, which depends on galaxy mass and redshift. The evolution of the stellar mass function for the two types is described by Muzzin et al. (2013), which we use here to compute this ratio. We also allow for 1% of outliers: these are objects that are not part of the galaxy population, for example, catastrophic redshift estimates or misclassified stars. Finally,

in order to avoid being dominated by the large number of low-mass galaxies, we also assign a weight to each galaxy which is inversely proportional to the number density. This ensures that each mass range carries equal weight in the fit. The number density is taken from the Muzzin et al. (2013) mass functions.

Then, we compute the total likelihood for a set of six model parameters (intercept A , slope α , and intrinsic scatter $\sigma_{\log R_{\text{eff}}}$, each for both types of galaxies):

$$\mathcal{L}_{\text{ET}} = \sum \ln \left[W \cdot \left((1 - C) \cdot P_{\text{ET}} + C \cdot P_{\text{LT}} + 0.01 \right) \right] \quad (5)$$

for early-type galaxies, and

$$\mathcal{L}_{\text{LT}} = \sum \ln \left[W \cdot \left((1 - C) \cdot P_{\text{LT}} + C \cdot P_{\text{ET}} + 0.01 \right) \right] \quad (6)$$

for late-type galaxies, where W is the weight and C is the contamination fraction, both of which are a function of redshift and mass. The best-fitting parameters are identified by finding the model with the maximum total likelihood, $\mathcal{L} = \mathcal{L}_{\text{ET}} + \mathcal{L}_{\text{LT}}$.

For the late types we fit all galaxies with $M_* > 3 \times 10^9 M_\odot$; this limit provides a good dynamic range of two orders of magnitude in mass, and exceeds the mass limit of our sample up to $z = 2.5$ (Figure 2). For the early types we fit all galaxies with $M_* > 2 \times 10^{10} M_\odot$, so that we avoid the clearly flatter part of the size-mass distribution at lower masses (see §3.2). This cut-off exceeds




Figure 6. Parametrized redshift evolution of the size-mass relation, from the power-law model fits shown in Figure 5. The left-hand panel shows the evolution of the intercept, or the size evolution at fixed stellar mass of $5 \times 10^{10} M_{\odot}$. Strong evolution is seen for high-mass early-type galaxies; moderate evolution for low-mass early types and for late-type galaxies. The middle and right-hand panels show the evolution of the slope and intrinsic (model) scatter of the size-mass relation, either with little or no evidence for changes with redshift. The open symbols represent the observed scatter: these measurements do not take measurement uncertainties and contamination into account. The fitting parameters shown in this figure are given in Table 1.




Figure 7. Evolution-corrected average sizes at $M_{*} = 5 \times 10^{10} M_{\odot}$ for late-type galaxies (top panel, in blue) and early-type galaxies (bottom panel, in red). The values shown here are the values shown in the left-hand panel of Figure 6, divided by $(1+z)^{\beta_z}$ as indicated on the y-axis. The residuals from the best-fitting $(1+z)^{\beta_z}$ law indicate that parametrizing the evolution as a function of the Hubble parameter ($R_{\text{eff}} \propto h(z)^{\beta_H}$) may provide a more accurate description for the late-type galaxies. See §3.2 for further discussion.

the mass limit of our sample up to $z = 3$.

The black lines in Figure 5 indicate the fitting results, and the evolution of the individual model parameters (intercept, slope and scatter) are shown in Figure 6. The fitting results are also given in Table 1. The intercept of the best-fitting size mass model distributions evolves significantly with redshift, and particularly rapidly for the early types.

Usually, the evolution of the intercept is parametrized as a function of $(1+z)$. While this is intuitively appealing because of our familiarity with the cosmologi-

cal scale factor, this is perhaps not the physically most meaningful approach. Galaxy sizes, in particular disk scale lengths, are more directly related to the properties of their dark matter halos than to the cosmological scale factor. Halo properties such as virial mass and radius follow the evolving expansion rate – the Hubble parameter $H(z)$ – instead of the cosmological scale factor. For a matter-dominated universe, $H(z)$ and $(1+z)$ evolve at a similar pace, but as a result of the increased importance at late times of Λ for the dynamical evolution of the universe, $H(z)$ evolves much slower in proportion to $(1+z)$ at late times than at early times. For example, at $z \sim 0$ we have $H(z) \propto (1+z)^{0.4}$, while at $z \sim 2$ this is $H(z) \propto (1+z)^{1.4}$.

For this reason it is reasonable to parametrize size evolution as a function of $H(z)$ in addition to $(1+z)$. The solid lines in the left-hand panel of Figure 6 represent the evolution as a function of $H(z)$, while the dashed lines represent the evolution as a function of $(1+z)$. These results are also given in Table 1. The $H(z)^{\beta_H}$ parametrization is marginally preferred by the data over the $(1+z)^{\beta_z}$ parametrization, as is more clearly illustrated in Figure 7, where we show the residuals. In addition to the statistical limitations, we note that these residuals are of the same magnitude as the systematic uncertainties in the size measurements and color gradient corrections (§2.5). A more thorough comparison with size evolution of larger samples at $z < 1$ with size measurements at visual wavelengths would improve these constraints.

Newman et al. (2012) first demonstrated the lack of strong evolution in the slope of the size-mass relation for massive ($> 2 \times 10^{10} M_{\odot}$) early-type galaxies. Here, we confirm that result (middle panel, Figure 6), and find a slope of $R_{\text{eff}} \propto M^{0.75}$ at all redshifts. This slope is somewhat steeper than measured by Shen et al. (2003) for present-day early-type galaxies. Differences in sample selection (star-formation activity vs. concentration) and methods (R_{eff} from Sérsic profile fits vs. Petrosian half-light radii) may explain this difference. For the first time we extend the analysis to late-type galaxies: the slope is much flatter than the slope for early types ($R_{\text{eff}} \propto M^{0.22}$), with little or no change with redshift.




Figure 8. Median (points) and 16th and 84th percentiles (lines) of the size-stellar mass distributions shown in Figure 5. The scatter in R_{eff} does not strongly depend on galaxy mass. Deviations from the power-law form of the size-mass relation are clear for massive late-type galaxies and for low-mass early-type galaxies. Note that here we do not account for contamination (misclassified early- and late-type galaxies).

This slope is intermediate to the slope found by Shen et al. (2003) for low- and high-mass galaxies. Our sample contains too few high-mass late-type galaxies to perform a robust double-component power-law fit, as done by Shen et al. (2003), but in §3.2 we will show evidence for the steepening of the relation for massive late-type galaxies out to $z = 1$.

Finally, we present the first measurement of the intrinsic scatter in size beyond the local universe (right-hand panel of Figure 6). We find no strong evolution for either late types or early types, and that the scatter for the early-type population is always somewhat smaller (0.1 to 0.15 dex) than for the late-type population (0.16 - 0.19 dex). These numbers agree well with the intrinsic scatter measured by Shen et al. (2003) for present-day galaxies: 0.13 dex for early-type galaxies, and 0.20 dex for late-type galaxies. We note that the effect of measurement uncertainties were not included by Shen et al. (2003).

For comparison we show the observed scatter at each redshift, calculated as the standard deviation in R_{eff} after subtracting the best-fitting size-mass relation. The values for early-type galaxies are in the range 0.2-0.3 dex, in good agreement with the values found by (Newman et al. 2012) over the same redshift range. In particular,

the strongly increased observed scatter in size for the early-type galaxies at $z > 2$ is largely attributed to significant contamination by mis-classified late-type galaxies. We have assumed a mis-classification probability of 10% (resulting in an assumed mis-classified fraction of $C = 0.10$ in the case of equal numbers of early- and late-type galaxies – see above), but although this value is empirically motivated it is not known with great precision. If we decrease (increase) the mis-classification probability to 5% (20%) then the recovered intrinsic scatter for the $z = 2.0 - 2.5$ early-type galaxy sample, for example, increases (decreases) to 0.18 (0.11).

At this point we should also comment on the effect of changing the value for the assumed random uncertainty in stellar mass (here, 0.15 dex). Decreasing its value has no measurable effect, while increasing it to 0.30 dex decreases the recovered value for the intrinsic scatter further, to 0.05 dex for the $z = 2.0 - 2.5$ early-type galaxy sample. In this sense, the derived values for the intrinsic scatter are upper limits.

While our particular choices in modeling the uncertainties affect the results with (marginal) significance, they do not affect our general conclusions that the intrinsic size scatter 1) is $\lesssim 0.20$ dex for both types of galaxies,




Figure 9. Median size as a function of stellar mass and redshift for early-type galaxies (left) and late-type galaxies (right). SDSS data points based on (Guo et al. 2009) are shown as open points. Fits to the median sizes of the form $R_{\text{eff}}/\text{kpc} = B_z(1+z)^{\beta_z}$ and $B_H(H(z)/H_0)^{\beta_H}$ are shown by dotted and solid lines, respectively. The evolution of the early-type galaxies is independent of mass at $M_* > 2 \times 10^{10} M_\odot$: massive galaxies evolve fast and have a steep size-mass relation at all redshifts, while the relation flattens out at lower masses ($\lesssim 10^{10} M_\odot$) and evolves less rapidly. The evolution of the late-type galaxies is overall slower, and does not depend strongly on mass. The low-mass early-type galaxies evolve at the roughly the same pace as the late-type galaxies. The median sizes and fitting results are given in Table 2.

and 2) does not strongly evolve with redshift. However, the conclusion that the scatter for early-type galaxies is smaller than for late-type galaxies at all redshifts – as is seen for present-day galaxies – should at this stage be regarded as tentative.

Finally, we note that changes in the mis-classification probability or uncertainty in stellar mass do not significantly affect the recovered values of the other model parameters (zero point and slope).

3.2. Evolution of median sizes

In this section we offer a complementary description of the evolution of the size-mass relation. In Figure 8 we show the median sizes as a function of mass and redshift, along with the 68th percentile width of the size distribution. The values are listed in Table 2. Up to $z \sim 1.5$ the relation for late types steepens and tightens at the high-mass end. Shen et al. (2003) modeled the steepening by assuming a two-component power law, but we sample an insufficiently large volume and sample size at $> 10^{11} M_\odot$ to include this in our analytical description presented above (§3.1). The flattening of the size-mass relation for low-mass early types is also clearly seen. Inspection of the SEDs of individual galaxies confirms that these are truly quiescent galaxies, with strong 4000Å breaks. As we showed in §3.1, the large apparent increase in the scatter for high-redshift early types can be partially attributed to contaminants and outliers.

We provide complementary sets of median size and scatter measurements in the Appendix. These include the commonly used circularized radii: $R_{\text{eff,circ}} = R_{\text{eff}}\sqrt{b/a}$, where b/a is the projected axis ratio. In addition, we provide the measurements for the combined late+early-type galaxy sample, and the measurements

in bins of rest-frame V-band luminosity.

Figure 9 shows the median size evolution for galaxies in different mass bins. We parametrize this evolution both as a function of $H(z)$ and of $(1+z)$ – see §3.1. The results are shown as solid and dotted lines, respectively, in Figure 9, and are also given in Table 2.

Ideally, an immediate comparison with the size-mass distribution of nearby galaxies provides a strong constraint on the evolution. However, such comparisons are fraught with systematic uncertainties. The aim here is merely to show that our observations from CANDELS and 3D-HST are consistent with the size-mass relation for nearby galaxies as measured from the SDSS (Shen et al. 2003), who provided the standard reference for this purpose.

In order to account for possible systematic differences we compare the size measurements from Shen et al. (2003) with those from Guo et al. (2009) on an object-by-object basis. The reason for using the Guo et al. (2009) measurements as a baseline is that they are based on the same technique – GALAPAGOS from Barden et al. (2012) – as used in this paper. We shift the analytical descriptions for the size-mass relations from Shen et al. (2003) according to the measured, systematic offset between Shen et al. (2003) and Guo et al. (2009). This amounts to a 0.1 dex shift to larger R_{eff} compared to Shen et al. (2003)¹⁹. In order to be conservative we also adopt 0.1 dex as the systematic uncertainty. We show the inferred sizes for local galaxies in Figure 9. The median size evolution traced out by the 3D-HST/CANDELS data predicts $z \sim 0$ sizes that are consistent with our adjusted Shen et al. (2003) median sizes.

¹⁹ Note that we use major axis R_{eff} in this paper, as opposed to Shen et al. (2003), who use circularized radii




Figure 10. Size distribution histograms for early- and late-type galaxies as a function of stellar mass (as labeled on the right-hand side) and redshift (as labeled at the top). The number of galaxies is given in units of co-moving volume to illustrate the growth of the population over time. The early-type size distributions are fit with Gaussians with a fixed dispersion of 0.16 dex. The late-type size distributions are fit with skewed Gaussians with a fixed dispersions of 0.18 dex and skewness $h_3 = -0.15$. The panels with thin lines show samples which are below our mass limit.




Figure 11. Cumulative size distributions of $\sim L^*$ early-type galaxies (top) and $\sim L^*$ late-type galaxies (bottom) as a function of redshift. While the number density of both early- and late-type galaxies increases over time, the number density of small galaxies declines, implying that the observed evolution in the mean size is not (solely) driven by the addition of larger galaxies. Individual galaxies must evolve in size.




Figure 12. Size evolution of galaxies in a narrow (0.3 dex) mass bin around $10^{10} M_{\odot}$. The black points represent UV-bright galaxies (with $U - V < 1$ in the rest-frame), selecting a sample akin to LBGs at high redshift. Their size evolution is fast, consistent with the size evolution of UV-selected samples up to $z = 7$ as recently determined by Mosleh et al. (2012) – also see Oesch et al. (2010). The blue points represent late-type galaxies as defined in this paper (see Figure 1), that is, all star-forming galaxies. The size evolution of those is slower at low redshift, consistent with previous measurements at $z < 1$ (here, Barden et al. 2005).

The picture provided by median size distributions is




Figure 13. Number density evolution of small, massive early-type galaxies. In each redshift bin (with equal co-moving volume) we include early-type galaxies with mass $M_* > 5 \times 10^{10} M_{\odot}$ and size $R_{\text{eff}}(\text{kpc}) < (M_*/5 \times 10^{10} M_{\odot})^{0.7}$. That is, the slope of the size-mass relation is taken into account: for $M_* = 5 \times 10^{10} M_{\odot}$ the size limit is $R_{\text{eff}} = 1$ kpc; for $M_* = 10^{11} M_{\odot}$ the size limit is $R_{\text{eff}} = 1.6$ kpc. The number density first increases with cosmic time, reaching a plateau at $z \sim 1.5 - 2$, after which it strongly decreases toward the present day. The immediate implication is that individual galaxies must grow in size significantly, most likely through merging.

consistent with our analytical description (§3.1), with fast evolution for the (massive) early types and moderate evolution for the late types. In addition, we see that the flattening of the relation for low-mass early types coincides with slower evolution. Interestingly, low-mass early-type galaxies evolve at a similar same rate as late types of the same mass. For the late types we see a mild dependence on mass: the more massive late types evolve slightly faster than the less massive late types. For further discussion, see §5.

3.3. Skewness in the R_{eff} distribution of late-type galaxies

The 16- and 84-percentile range for late-type galaxies is not precisely centered on the median size (Figure 8 and Table 2), implying a skewness in the size distribution. To examine this further we show size distribution histograms for a set of mass and redshift bins in Figure 10. The asymmetric size distribution for late-type galaxies is due to a tail of small galaxies. The small sizes are not due to point-like contributions from AGN: the $1 - 10 \mu\text{m}$ photometry of these objects does not show the power-law spectral energy distributions that are characteristic for unobscured AGN.

As a result of this skewness there is substantial overlap between the size distributions of early types and late types and no clear bi-modality; this, despite large differences in the average sizes. Figure 11 shows – more clearly than Figure 10 – that the size distributions of the two types overlap at all redshifts, up to $z \sim 3$.

Table 1

Results from the parametrized fits to the size-mass distribution of the form $R_{\text{eff}}/\text{kpc} = A(M_*/5 \cdot 10^{10} \text{ M}_\odot)^\alpha$, as described in §3.1 and shown in Figures 5 and 6. $\sigma(\log R_{\text{eff}})$ is the scatter in R_{eff} in logarithmic units.

z	Early-Type Galaxies			Late-Type Galaxies		
	$\log(A)$	α	$\sigma \log(R_{\text{eff}})$	A	α	$\sigma \log(R_{\text{eff}})$
0.25	0.60 ± 0.02	0.75 ± 0.06	0.10 ± 0.02	0.86 ± 0.02	0.25 ± 0.02	0.16 ± 0.01
0.75	0.42 ± 0.01	0.71 ± 0.03	0.11 ± 0.01	0.78 ± 0.01	0.22 ± 0.01	0.16 ± 0.01
1.25	0.22 ± 0.01	0.76 ± 0.04	0.12 ± 0.01	0.70 ± 0.01	0.22 ± 0.01	0.17 ± 0.01
1.75	0.09 ± 0.01	0.76 ± 0.04	0.14 ± 0.01	0.65 ± 0.01	0.23 ± 0.01	0.18 ± 0.01
2.25	-0.05 ± 0.02	0.76 ± 0.04	0.14 ± 0.02	0.55 ± 0.01	0.22 ± 0.01	0.19 ± 0.01
2.75	-0.06 ± 0.03	0.79 ± 0.07	0.14 ± 0.03	0.51 ± 0.01	0.18 ± 0.02	0.19 ± 0.01

Table 2

Logarithmic size distribution (16% – 84% range) as a function of galaxy mass and redshift. The masses in the header and the redshifts in the first column are the centers of 0.5-wide bins. Redshift dependence in the form $R_{\text{eff}}/\text{kpc} = B_z(1+z)^{\beta_z}$ and $R_{\text{eff}}/\text{kpc} = B_H(H(z)/H_0)^{\beta_H}$ are also given.

z	Early-Type Galaxies					Late-Type Galaxies				
	$M_* = 10^{9.25}$	9.75	10.25	10.75	11.25	9.25	9.75	10.25	10.75	11.25
0.25	16%	0.03 ± 0.06	0.04 ± 0.03	0.13 ± 0.03	0.42 ± 0.02	0.65 ± 0.05	0.24 ± 0.01	0.36 ± 0.02	0.42 ± 0.02	0.61 ± 0.03
	50%	0.27 ± 0.02	0.28 ± 0.02	0.38 ± 0.03	0.67 ± 0.02	0.76 ± 0.09	0.49 ± 0.01	0.61 ± 0.02	0.66 ± 0.03	0.83 ± 0.03
	84%	0.46 ± 0.03	0.46 ± 0.04	0.58 ± 0.04	0.92 ± 0.05	1.08 ± 0.08	0.70 ± 0.01	0.80 ± 0.01	0.85 ± 0.02	1.01 ± 0.04
0.75	16%	-0.02 ± 0.03	-0.14 ± 0.02	0.02 ± 0.02	0.26 ± 0.01	0.62 ± 0.02	0.18 ± 0.01	0.32 ± 0.01	0.39 ± 0.01	0.51 ± 0.02
	50%	0.23 ± 0.01	0.21 ± 0.02	0.23 ± 0.01	0.45 ± 0.01	0.81 ± 0.02	0.43 ± 0.01	0.56 ± 0.01	0.64 ± 0.01	0.75 ± 0.01
	84%	0.43 ± 0.02	0.44 ± 0.02	0.42 ± 0.02	0.64 ± 0.02	0.97 ± 0.02	0.65 ± 0.01	0.76 ± 0.01	0.83 ± 0.01	0.90 ± 0.01
1.25	16%	...	-0.15 ± 0.03	-0.15 ± 0.02	0.07 ± 0.01	0.41 ± 0.02	0.11 ± 0.01	0.23 ± 0.01	0.33 ± 0.01	0.47 ± 0.01
	50%	...	0.18 ± 0.03	0.09 ± 0.02	0.30 ± 0.01	0.58 ± 0.03	0.37 ± 0.01	0.48 ± 0.01	0.57 ± 0.01	0.67 ± 0.01
	84%	...	0.42 ± 0.04	0.36 ± 0.03	0.54 ± 0.03	0.81 ± 0.03	0.60 ± 0.01	0.69 ± 0.00	0.77 ± 0.01	0.83 ± 0.01
1.75	16%	...	-0.02 ± 0.06	-0.27 ± 0.02	-0.04 ± 0.02	0.28 ± 0.02	0.07 ± 0.01	0.16 ± 0.01	0.28 ± 0.01	0.35 ± 0.02
	50%	...	0.22 ± 0.03	0.02 ± 0.03	0.19 ± 0.02	0.45 ± 0.02	0.33 ± 0.01	0.42 ± 0.01	0.52 ± 0.01	0.61 ± 0.01
	84%	...	0.48 ± 0.06	0.35 ± 0.03	0.50 ± 0.03	0.74 ± 0.04	0.57 ± 0.01	0.65 ± 0.01	0.72 ± 0.01	0.80 ± 0.01
2.25	16%	-0.37 ± 0.08	-0.20 ± 0.02	0.16 ± 0.03	...	0.10 ± 0.01	0.17 ± 0.02	0.26 ± 0.03
	50%	-0.04 ± 0.07	0.08 ± 0.03	0.36 ± 0.05	...	0.35 ± 0.01	0.44 ± 0.01	0.53 ± 0.01
	84%	0.36 ± 0.03	0.54 ± 0.04	0.55 ± 0.07	...	0.57 ± 0.01	0.64 ± 0.01	0.70 ± 0.01
2.75	16%	-0.22 ± 0.05	0.07 ± 0.07	0.16 ± 0.02	0.19 ± 0.06
	50%	0.10 ± 0.03	0.39 ± 0.08	0.43 ± 0.01	0.47 ± 0.01
	84%	0.50 ± 0.10	0.68 ± 0.10	0.65 ± 0.02	0.71 ± 0.02
B_z	...	0.29 ± 0.01	0.47 ± 0.02	0.75 ± 0.04	1.05 ± 0.09	0.54 ± 0.01	0.69 ± 0.01	0.74 ± 0.03	0.90 ± 0.05	1.05 ± 0.08
β_z	...	-0.22 ± 0.02	-1.01 ± 0.06	-1.24 ± 0.08	-1.32 ± 0.21	-0.48 ± 0.03	-0.63 ± 0.02	-0.52 ± 0.08	-0.72 ± 0.09	-0.80 ± 0.18
B_H	...	0.27 ± 0.01	0.42 ± 0.02	0.68 ± 0.03	0.97 ± 0.06	0.52 ± 0.01	0.65 ± 0.01	0.71 ± 0.03	0.86 ± 0.04	1.01 ± 0.06
β_H	...	-0.19 ± 0.02	-0.97 ± 0.05	-1.13 ± 0.06	-1.29 ± 0.16	-0.52 ± 0.02	-0.58 ± 0.02	-0.49 ± 0.07	-0.65 ± 0.09	-0.76 ± 0.13

Figure 10 also shows tails of large, early-type galaxies. However, this can likely be attributed to misclassification: their number is always an order magnitude smaller than the number of late-type galaxies with similar sizes, consistent with the assumed mis-classification probability in our analysis in Section 3.1. We note that the tail of small, late-type galaxies is not consistent with the expected number of mis-classified objects: its prominence appears to be unrelated to the early-type population.

As an illustration that the size distribution for both early- and late-type galaxies evolve smoothly and regularly, we fit Hermite-Gaussian functions to the histograms shown in Figure 10. This provides a reasonable description for all redshift and mass bins.

For a discussion of the implications of these results in the context of previous results we refer to §4.2 and §5.2.

4. COMPARISON WITH PREVIOUS RESULTS

4.1. Late-Type Galaxies

The results for size evolution of late-type galaxies shown in Figures 6 and 9 are consistent with most other recent measurements that focus on the $z = 1 - 2$ redshift range (e.g., Franx et al. 2008; Buitrago et al. 2008). Williams et al. (2010) found somewhat faster evolution for late types (and slower evolution for early types), but a direct comparison with their Figure 4 reveals that their and our size measurements are in fact fully consistent over the redshift range $0.5 < z < 1.5$. The difference in the quoted pace of evolution is likely the result of the increased dynamic range in redshift probed here, in addition to the use of a low-redshift comparison sample from the SDSS (see §3.2).

Thus, like previous studies focusing on the $z = 1 - 2$ redshift regime, we find that the pace of evolution for late-type galaxies is intermediate to the slow evolution of disk galaxies at $z < 1$ (Lilly et al. 1998; Ravindranath et al. 2004; Barden et al. 2005) and the fast evolution of UV-selected galaxies at $z > 2$ (Giavalisco et al. 1996; Ferguson et al. 2004; Oesch et al. 2010; Mosleh et al. 2012). Our data set allows us to bridge these regimes and probe the origin of this difference.

In Figure 12 we show the size evolution of galaxies with stellar mass $M_* \sim 10^{10} M_\odot$ from the present day up to $z \sim 6$. Here we have relaxed our magnitude limit to $H_{F160W} = 26$, which is still within the completeness limit of the CANDELS imaging as can be seen in Figure 2. Size measurements of individual galaxies are no longer reliable at $H_{F160W} = 26$, but the sample average is still robust to within 15% (van der Wel et al. 2012). For galaxies bluer than $U - V = 1$, thus selecting a population akin to LBGs we probe the population out to $z \sim 6$. The median size evolves quickly with redshift, $R_{\text{eff}} \propto (1 + z)^{-1.1}$, consistent with recent measurements by Oesch et al. (2010) and Mosleh et al. (2012).

Once we include all late-type galaxies, regardless of color, the evolution matches that of the $U - V < 1$ galaxies at $z \gtrsim 2$. This is simply because essentially all galaxies are blue: the color-blind sample is not biased against galaxies with $U - V > 1$ up to $z = 3 - 3.5$. At lower redshift red star-forming galaxies appear, which are smaller in size than their blue counterparts and slow down the average size evolution. At $z < 1$ UV-bright galaxies are

rare and the evolution is dominated by redder galaxies, which evolve in size more slowly in agreement with the results from, for example, Barden et al. (2005).

As we argued in §3.2 galaxy sizes may be better parametrized as a function of $H(z)$ than as a function of $(1 + z)$. While the former naturally implies slower evolution at late times than the latter (see the red, dotted line in Figure 12) not all trends are captured by using the $H(z)$ parametrization: 1) the evolution of all late types slows down more rapidly than can be explained by the difference between the two parametrizations; 2) the UV-bright sample shows little evidence for slowed evolution at $z \lesssim 1$.

We conclude that the diverging pace of evolution seen at $z < 1$ and $z > 3$ as reported in the literature is in part due to sample selection effects and part due to the different evolution of red and blue late-type galaxies.

4.2. Early-Type Galaxies

As we discussed in the Introduction there is broad agreement that the average sizes of $\sim L^*$ early-type galaxies, as measured at a fixed stellar mass, were smaller at high redshift. Moderate deviations in the pace of evolution can be attributed to sample selection-, measurement- and/or fitting techniques. For example, as we mentioned above, whereas the size measurements reported by Williams et al. (2010) are consistent with our size measurements, the reported pace of evolution is somewhat different: $(1 + z)^{-1}$ from Williams et al. (2010); $(1 + z)^{-1.3}$ from §3.2 in this paper. This is the result of the difference in spanned redshift range and the different use of present-day comparison samples. While these differences are large enough to be interesting, there is reasonable consensus that the average size for the population of early-type galaxies evolves rapidly. In particular, the first HST/NICMOS-based produced an impressive body of evidence for rapid evolution (Zirm 2007; Toft et al. 2007; van Dokkum et al. 2008; Buitrago et al. 2008), later confirmed by HST/WFC3-based studies (e.g., Newman et al. 2012; Cassata et al. 2013).

What has so far remained contentious is what drives this evolution: size evolution of individual galaxies, the addition of larger galaxies to the population, or a combination. Figure 11 (top panel) shows the cumulative size histograms of early-type galaxies in the L^* mass range, which reveals that the number density of small early-type galaxies strongly evolves with redshift. The total number density of early-type galaxies increases from $z \sim 3$ to the present, but the number of small galaxies strongly decreases at late cosmic times. We show this explicitly in Figure 13: the number density of high-mass galaxies with small sizes increases from early times to $z \sim 2$, levels off, and then decreases strongly at $z \lesssim 1.5$. The definition of small is arbitrary, but the generally picture does not depend on the precise choice in mass and size range. This finding is in general agreement with previous claims based on smaller samples and fewer fields by Cassata et al. (2011), Newman et al. (2012), Szomoru et al. (2012), Buitrago et al. (2013), and Cassata et al. (2013).

Several authors have argued that there are substantial numbers of small, yet massive galaxies in the present-day universe (Valentinuzzi et al. 2010; Poggianti et al. 2013). The latter show that 3 – 5% of present-day group and cluster early-type galaxies with mass $M_* > 3 \times 10^{10} M_\odot$

can be classified as “compact”. Following their definition ($M_*/(2\pi R_{\text{eff,circ}}) > 3 \times 10^9 M_\odot \text{ kpc}^{-2}$) we find that $\sim 40 - 50\%$ of $z \sim 1$ early-type galaxies qualify as compact. Since the total number density of such galaxies has evolved by no more than a factor 2 or 3 we conclude that most of the $z \sim 1$ compact galaxies no longer exist in that form in the present-day universe. Several ‘fossils’ in the form of $R_{\text{eff}} \sim 1 \text{ kpc}$, $M_* \sim 10^{11} M_\odot$ galaxies are found in the local volume (see, e.g., van den Bosch et al. 2012; Dullo & Graham 2013, for recent examples), but their number density is too low to match the number density of their $z = 2$ counterparts.

Recently Carollo et al. (2013) claimed that the number density of small early-type galaxies in the L^* mass range has not strongly evolved since $z \sim 1$. We rule out that field-to-field variations explain the discrepancy with our results. All five fields show a decline in the number density of compact galaxies (as defined in Figure 13, with $R_{\text{eff}} < 2.5$ at $10^{11} M_\odot$) between $z = 1.5$ and $z = 0.5$, by factors ranging from 3 to 10. A decline of more than a factor 2 between $z = 1$ and $z = 0.5$ is seen for 4 out of 5 fields.

There are several factors, in the form of systematic effects in the size and mass measurements, that may contribute to this tension. Slight redshift-dependent shifts in the stellar mass estimates produce changes in the size distribution as measured at a fixed mass. Our stellar mass estimates for luminous early-type galaxies have been demonstrated to show small, if any, shifts ($\lesssim 0.1$ dex) compared to dynamical mass measurements over the redshift range $0 < z < 2$ (van de Sande et al. 2013; Bezanson et al. 2013; Belli et al. 2014). In addition, our color-gradient correction would introduce a 14% shift of $z = 1$ sizes relative to $z = 0.2$ sizes in the Carollo et al. sample.

Most importantly, the size measurements used here and by Carollo et al. (2013) are obtained with fundamentally different techniques: here we use parametrized profile fits, while Carollo et al. use growth curves. The growth-curve method does not take the PSF into account at the time of measurement, but relies on a *a posteriori* correction. The magnitude of the correction depends on the intrinsic structural properties and is inferred from simulated size measurements. For example, galaxies with measured sizes of $\sim 0.2''$ receive a negligible correction, whereas galaxies with measured sizes of $\sim 0.1''$ receive a factor 2 correction downward (see Figure 2 of Carollo et al.). With such strongly size-dependent corrections it is difficult to reconstruct the true size distribution, especially when those corrections are of similar magnitude as the sizes themselves.

In an explicit example in which we apply a systematic, redshift-dependent shift in R_{eff} of order 0.1-0.2 dex per unit redshift we infer non-evolution in the number density of compact galaxies. Given this sensitivity to small shifts in size we argue that our measurements, which do not require systematic size corrections of more than a few percent (see Section 2.5 and van der Wel et al. (2012)), represent the size distribution with good fidelity across the examined redshift range.

5. DISCUSSION

5.1. Evolution of Late-Type Galaxies

Remarkably, the observed pace of size evolution for late-type galaxies is essentially the same as the evolution of the dark matter halo radius at a fixed mass, $R \propto H(z)^{-2/3}$, but only when halo mass and radius are defined with respect to the critical density. In a Λ CDM universe, if halo mass is parametrized with respect to matter density or virial density (assuming top-hat collapse), then Λ causes strong departures from a power law at late cosmic times. The average evolution between $z = 2$ and the present is $R \propto H(z)^{-1.06}$ or $\propto H(z)^{-1.24}$, respectively (Peebles 1980).

The interpretation of such a comparison is not straightforward. However, our novel measurement of the slope and scatter of the size mass relation provides new constraints. The intrinsic scatter in galaxy size remains approximately the same at all redshifts ($\sim 0.16 - 0.19$ dex, see Figure 6) and is comparable to, but perhaps somewhat smaller than, to the scatter of 0.25 dex in the halo spin parameter (e.g., Macciò et al. 2008). This strongly suggests that at all redshifts the sizes of late-type galaxies are set by their dark matter halos, and encourage us to examine the relation between galaxy sizes and halo properties further.

The power-law fits presented in §3.1 imply that there is very little or no evolution in the slope of the size-mass relation; it remains flat, $\alpha \equiv d \log R_{\text{eff}} / d \log M_* = 0.22 \pm 0.03$, at all redshifts $0 < z < 3$ (Figure 6, middle panel). As argued by Shen et al. (2003), the flat slope suggests that the ratio between galaxy mass and halo mass is not a constant: if it were, the size-mass relation would be steeper ($\alpha = 1/3$) than observed. The underlying assumption is that galaxy size is proportional to halo size (Kravtsov 2013), which we here take to be the case for late-type galaxies. Following Shen et al. (2003), we use the observed slope ($\alpha \sim 1/5$) to constrain the galaxy mass-halo mass relation and find $m_g \equiv M_{\text{gal}}/M_{\text{halo}} \propto M_{\text{halo}}^{\gamma \sim 2/3}$. The observation that the slope of the size-mass relation does not evolve with redshift provides a very stringent constraint on the models: unless a combination of factors conspire to keep this slope constant, the most straightforward explanation is that the slope of the relation between galaxy and halo mass (γ) is similar across the redshift range considered here.

Indeed, entirely independent estimates of the relationship between galaxy and halo mass, based on clustering measurements and abundance matching techniques, provide strong evidence that m_g depends on halo mass, similarly so at different redshifts (e.g., Conroy & Wechsler 2009; Moster et al. 2010; Behroozi et al. 2010; Wake et al. 2011; Moster et al. 2013; Behroozi et al. 2013a). In fact, the most recent studies found that $\gamma = 2/3$ for halos in the mass range $M_{\text{halo}} \sim 10^{11-12} M_\odot$, in agreement with what we infer based on the slope of the size-mass relation. In addition, Moster et al. (2013) and Behroozi et al. (2013a) showed that m_g peaks at a similar halo mass ($\sim 10^{12} M_\odot$) at all redshifts $z \lesssim 2$, at around a constant value of $m_g \sim 0.05$. The implication is that m_g does not strongly evolve over the (rather narrow) halo mass range $10^{11-12} M_\odot$ (Behroozi et al. 2013b).

It is unclear whether the observed pace of galaxy size evolution ($R_{\text{eff}} \propto H(z)^{-2/3}$) implies that R_{eff} evolves in proportion to R_{halo} as may be expected in the case that galaxy size scales with halo size in the present-day

universe (Kravtsov 2013): it may be a coincidence that the observed pace of evolution is the same as that for halo radii with respect to the critical density: it appears more natural to expect galaxy sizes to scale with halo mass and radius that are defined in terms of matter or virial density. In this spirit, the tendency for late-type galaxies to display rather slower size evolution than expected has been given ample attention in the literature.

Somerville et al. (2008) argued that because halos are less concentrated at high redshift baryonic disks are larger in proportion to the virial radii of halos, leading to slower size evolution. In addition, Dutton et al. (2011) showed that accreting, gaseous disks with a simple, but self-consistent prescription for star formation lead to similarly slow evolution of the stellar disk scale radius as a result of recycling of gas and radial variations in star formation. In addition, stellar feedback may have a more direct impact on disk sizes as low-angular momentum material is ejected (e.g., Maller & Dekel 2002; Brook et al. 2011).

The sizes we measured are the not strictly disk scale lengths, as we sample the whole galaxy, including the bulge. Therefore, bulge formation in late-type galaxies slows down size evolution as parametrized here. Bulge formation can either occur rapidly, through mergers (Toomre & Toomre 1972) or clump formation and migration in unstable disks (Dekel et al. 2009; Ceverino et al. 2010), or gradually, through secular evolution driven by non-axisymmetries in the disk potential (Kormendy & Kennicutt 2004). The prediction of any of these scenarios is that the galaxies with higher global Sérsic indices will have smaller sizes at a given mass. The observation that evolution is faster at $z > 2$ and slower at $z < 1$ (see §4.1 and Figure 12), combined with the appearance of redder, more compact galaxies at late cosmic times, suggests that bulge formation plays an important role in the evolution of half-light radii of late-type galaxies.

5.2. Evolution of Massive Early-Type Galaxies

The co-moving number density of L^* early types has strongly increased over the redshift range examined in this paper ($0 < z < 3$), as was shown by, e.g., Bell et al. (2004); Faber et al. (2007); Brown et al. (2007); Ilbert et al. (2010); Brammer et al. (2011); Buitrago et al. (2013); Muzzin et al. (2013). Here, this is illustrated in Figures 10 and 11. Naturally, the progenitors of the newly formed early-type galaxies must be looked for among the star-forming, late-type population. The skewed size distribution of late types toward small sizes (see §3.3 and Figure 10) points at the existence of a population of small late-type galaxies that span the entire size range seen for early-type galaxies. Figure 11 illustrates that this is the case over essentially the entire redshift range probed by our sample.

The tail of small, star-forming galaxies shown in Figure 11 at $z > 1.5$ (also see Barro et al. 2013; Williams et al. 2014; Barro et al. 2013) may reflect the intriguing possibility of a scenario in which such small, yet massive star-forming galaxies are the immediate progenitors of compact early-type galaxies. Their number density does not rapidly change between $z = 3$ and $z = 1.5$, whereas the number of early-type galaxies does rapidly increase over that redshift range (see Figure 11). This would suggest the continuous emergence of additional, small late-

type galaxies that represent a transitional phase between the bulk of the late-type population and the early-type population, as recently advocated by Dekel & Burkert (2014) based on analytical arguments and simulations.

An alternative interpretation is that the star-forming population consists of ‘normal’ late types and a population of early-type galaxies that revived their star-formation activity. The simplest implementation of this model, in which these ‘frosting’ early types have the same size distribution of the quiescent early types, can be ruled out: the skewed size distribution of late types is not well described by two log-normal distributions centered at the respective peaks of the size distributions for late- and early-type galaxies. In general, the size distribution of the full population of galaxies (early and late types combined) is not observed to be bimodal in the sense that there is no clear gap between two fiducial populations of small and large galaxies; nor can the size distribution be accurately represented by a single Gaussian distribution. More complicated models of the ‘frosting’ flavor, in which a large, star-forming disk re-assembles to surround a compact, quiescent component cannot be immediately ruled out. However, such scenarios seem implausible as the implied color and mass-to-light ratio gradients of such galaxies would likely be stronger than observed (Wuyts et al. 2012). Measurements of the stellar density in the central regions of early- and late-type galaxies can be used to provide further constraints.

Whether or not the small late-type galaxies represent a transitional phase, the central idea in the formation of early-type galaxies is that the formation of an early-type galaxy occurs requires the formation of a concentrated stellar body with a high density (e.g., Franx et al. 2008; Bell et al. 2012). One possibility is that a substantial amount of material flows to the center under the influence of mergers (e.g., Di Matteo et al. 2005) or violently unstable disks and clump formation/migration (Dekel et al. 2009; Ceverino et al. 2010; Dekel & Burkert 2014). It remains to be seen whether such processes can reproduce the correct stellar density profiles (Wuyts et al. 2010).

As we showed in §4.2 and Figures 11 and 13 the number density of small, compact early-type galaxies strongly decreases between $z \sim 1.5$ and the present. This immediately implies that early-type galaxies, after they first form as compact, quiescent objects, have to substantially grow in size over time. Combining this with the suggestion that new early-type galaxies likely form out of the smallest late-type galaxies, the implication is that early-type galaxies are the most dense (and disk-like in structure, e.g., van der Wel et al. 2011; Bruce et al. 2012; Chang et al. 2013) immediately after their star formation is truncated. The amount of later evolution in size and density is dictated by the (non-evolving) slope of the size-mass relation and the evolution of its intercept. This naturally fits into the general idea that a gas-rich formation phase is followed by a more quiescent, dissipationless formation phase.

The scatter in the size-mass relation of $\sim 0.15 - 0.20$ dex (see Figure 6) shows that there is some variation in amount of dissipative and dissipationless formation, yet, the fact that we see little or no evolution in the size scatter implies that the amount of dissipation integrated over cosmic history does not vary greatly; some early-type galaxies may have experienced an intensely dissi-

pative phase at early times, while other, but similarly massive, galaxies may have continued a less intense star-forming phase up until recently. The compact $z > 1.5$ early-type galaxies would fall in the former category; the large, massive star-forming galaxies at $z \sim 0.5 - 1.5$ may be the progenitors of galaxies in the latter.

Within this framework, independent evidence for the increase in stellar mass of individual early-type galaxies by a factor of 2 – 3 between $z = 2$ and the present (van Dokkum et al. 2010) implies that the growth in size depends on the growth in mass as $\Delta R_{\text{eff}} \propto \Delta M_*^{0.2}$. This steep dependence is consistent with a merger scenario.

Satellite galaxies can be stripped and their stars deposited on large-radius orbits. Direct and stringent constraints on the minor merger rate are difficult to obtain, but it has proved to be difficult to observationally confirm a sufficiently large minor merger rate to explain the observed evolution (Newman et al. 2012). Mergers among galaxies that occupy the size-mass relation for early-type galaxies, that is, pure dry mergers, may not occur at sufficient rates (e.g., Nipoti et al. 2012).

Alternatively, mergers between similarly massive galaxies with different sizes can induce large changes in the size-mass distribution of the population. Assuming that the size distribution of progenitors partaking in major mergers is the same as that of the population as a whole, a $R_{\text{eff}} = 1$ kpc early-type galaxy at $z \sim 2$ will merge, on average, with a late-type galaxy that is 3 times larger. The properties of the merger remnant will depend on the amount of dissipation and the dynamics of the merger, but it is conceivable that the remnant will be much larger than the compact progenitor. A dense inner region will remain in place, and the strong correlation between central density and quiescence implies that the remnant is likely to be quiescent as well.

The mass ratio distribution in the merger history of early-type galaxies, and its effect on size evolution, will remain a topic of debate. However, merging can account for, and is arguably required to explain, the disappearance from $z \sim 2$ to the present of disk-like structures among L^* early types (van der Wel et al. 2011; Bruce et al. 2012; Chang et al. 2013), and the observation that the most massive galaxies in the present-day universe do not have a disk-like structure, but are intrinsically round (van der Wel et al. 2009). A combined analysis of the evolution of size and morphological properties (see, e.g., Huertas-Company et al. 2013) will aid to simultaneously interpret size growth and disk destruction.

The above narrative shows that we have gathered a plausible set of mechanisms that may play a role in explaining the formation and subsequent evolution of early-type galaxies. Despite this we lack the basis of a simple analytical framework that is similar to our model for disk formation. However, we note that the rapid pace of size evolution is very close to the size evolution expected for halos as defined by their virial density: $R \propto H(z)^{-1.24}$ for halos compares well with $R_{\text{eff}} \propto H(z)^{-1.29}$ for massive early-type galaxies. If we assume that these galaxies only grow through the accretion of other halos and their stellar content, then it is perhaps not a coincidence that halos and galaxies both follow the evolutionary path expected for a dissipationless, top-hat collapse scenario.

5.3. Evolution of Low-Mass Early-Type Galaxies

As we noted in §3.2 the slope of the size-mass relation for early-type galaxies flattens below stellar mass $\sim 10^{10} M_\odot$, and the size evolution is more comparable to that of late types than that of early types (see Figure 9). This suggests that there is a population of low-mass early types that may have formed out of late-type galaxies without going through a transitional phase in which high central densities are attained. Stripping of gas from satellite galaxies is a natural explanation for such evolution, and can explain the existence of an excess population of early-type galaxies in clusters that have structural properties similar to those of late-type galaxies in the field (van der Wel et al. 2010). Satellites are common in this mass range in the present-day universe (e.g., van den Bosch et al. 2008), but not at higher redshifts, lending the stripping scenario more credence based on the rapid increase in the co-moving number density since $z \sim 1 - 1.5$ (see Figures 5 and 10).

On the other hand, the early types with mass $\lesssim 10^{10} M_\odot$ are ~ 2 times smaller than equally massive late types. Disk fading may contribute to this difference, but bulge formation and, in general, processes that cause more massive galaxies to transform into early-type galaxies, may play a role in the low-mass regime as well. A model such as that presented by Peng et al. (2010) can be expanded in order to separately reproduce the size-mass relations for different types of ‘quenched’ galaxies.

6. SUMMARY

In this paper we present the size-mass distribution of 30,958 galaxies over a large range in mass ($> 10^9 M_\odot$) and redshift ($0 < z < 3$), distinguishing between early-type and late-type galaxies based on their star-formation activity. Spectroscopic and photometric redshifts, stellar masses and rest-frame properties are determined using data from the 3D-HST survey and auxiliary, multi-wavelength photometric data sets spanning from the U band to $8\mu\text{m}$ (see §2). Galaxy sizes are measured from CANDELS imaging by single-component Sérsic profile fits to two-dimensional light distributions, with a correction for (redshift-dependent) color gradients (§2.5).

Consistent with previous results, we find that high-redshift galaxies are substantially smaller than equally massive, present-day counterparts. As is shown in Figures 5, 6 and 9, late-type galaxies are, on average, a factor ~ 2 smaller at $z = 2$ than at the present-day, whereas for massive early-type galaxies this is a factor of ~ 4 . We find that the size evolution of late-type galaxies is marginally better described as a function of the redshift-dependent Hubble parameter, $H(z)$, than as a function of the scale factor, $1 + z$ (Figure 7). Average, mass-matched sizes of late- and early-type galaxies evolve as $R_{\text{eff}} \propto H(z)^{-0.66} \propto (1 + z)^{-0.75}$ and $R_{\text{eff}} \propto H(z)^{-1.29} \propto (1 + z)^{-1.48}$, respectively (Figure 6 and Table 1).

High-mass late-type galaxies evolve marginally faster than low-mass late-type galaxies (Figure 9 and Table 2), but the data are consistent with no evolution in the overall slope of the size-mass relation. At all redshifts $z \leq 3$ we find that the slope is shallow for late-type galaxies ($R_{\text{eff}} \propto M_*^{0.22}$ for galaxies with stellar mass $M_* > 3 \times 10^9 M_\odot$), and steep for early-type

galaxies ($R_{\text{eff}} \propto M_*^{0.75}$ for galaxies with stellar mass $M_* > 2 \times 10^{10} M_\odot$). The size-mass relation for lower-mass early-type galaxies is more similar to that of late types than that of high-mass early types (§5.3). Once cross-contamination between the two classes of galaxies and outliers are taken into account (Figure 6 and §3.1), we also find no evidence for evolution in the (intrinsic) size scatter at a fixed galaxy mass. The implications of these results are discussed in §5.

The data presented here are consistent with essentially most published data sets (§4). Because of the sample size and dynamic range in mass and redshift, the immediate implications of the measurements are less ambiguous than was the case for previous studies. In particular, we show in Figure 11 that the size distribution of $z \sim 2$ early-type galaxies is significantly different from that of any subset of low-redshift galaxies with the same comoving number density; small early-type galaxies, which are typical at $z \sim 2$, do not exist in equal numbers today (Figure 13) and must therefore undergo strong size evolution in the intervening time.

The size-mass distributions from the 3D-HST and CANDELS projects presented here provide a solid framework for galaxy evolution models. The steadily evolving intercept of the size-mass relation, in combination with the non-evolving slope and scatter, present tight constraints on how baryons condense and form galaxies at the centers of dark matter halos (e.g., §5.1). The different assembly mechanisms of early- and late-type galaxies act similarly at all redshifts, as evidenced by the very different, but unchanging slopes of their respective size-mass relations.

This work is based on observations taken by the CANDELS Multi-Cycle Treasury Program (PI: Faber) and the 3D-HST Treasury Program (PI: van Dokkum) with the NASA/ESA HST, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

REFERENCES

- Barden, M. et al. 2005, *ApJ*, 635, 959
 Barden, M., Häußler, B., Peng, C. Y., McIntosh, D. H., & Guo, Y. 2012, *MNRAS*, 422, 449
 Barro, G. et al. 2013, *ApJ*, 765, 104
 Barro, G. et al. 2013, arXiv:1311.5559
 Behroozi, P. S., Conroy, C., & Wechsler, R. H. 2010, *ApJ*, 717, 379
 Behroozi, P. S., Wechsler, R. H., & Conroy, C. 2013, *ApJ*, 770, 57
 Behroozi, P. S., Wechsler, R. H., & Conroy, C. 2013, *ApJ*, 762, L31
 Bell, E. F. et al. 2004, *ApJ*, 608, 752
 Bell, E. F., van der Wel, A., Papovich, C., et al. 2012, *ApJ*, 753, 167
 Belli, S., Newman, A. B., & Ellis, R. S. 2014, *ApJ*, 783, 117
 Bezanson, R., van Dokkum, P. G., Tal, T., Marchesini, D., Kriek, M., Franx, M., & Coppi, P. 2009, *ApJ*, 697, 1290
 Bezanson, R., van Dokkum, P. G., van de Sande, J., et al. 2013, *ApJ*, 779, L21
 Bédorf, J., & Portegies Zwart, S. 2013, *MNRAS*, 867
 Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *ApJ*, 686, 1503
 Brammer, G. B. et al. 2011, *ApJ*, 739, 24
 Brammer, G. B. et al. 2012b, *ApJ*, 758, L17
 Brammer, G. B. et al. 2012a, *ApJS*, 200, 13
 Brown, M. J. I. et al. 2007, *ApJ*, 654, 858
 Brook, C. B. et al. 2011, *MNRAS*, 415, 1051
 Bruce, V. A. et al. 2012, *MNRAS*, 427, 1666
 Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
 Buitrago, F., Trujillo, I., Conselice, C. J., Bouwens, R. J., Dickinson, M., & Yan, H. 2008, *ApJ*, 687, L61
 Buitrago, F., Trujillo, I., Conselice, C. J., Häußler, B. 2013, *MNRAS*, 428, 1460
 Carollo, C. M. et al. 2013, *ApJ*, 773, 112
 Cassata, P. et al. 2011, *ApJ*, 743, 96
 Cassata, P. et al. 2013, *ApJ*, 775, 106
 Ceverino, D., Dekel, A., & Bournaud, F. 2010, *MNRAS*, 404, 2151
 Chabrier, G. 2003, *PASP*, 115, 763
 Chang, Y.-Y. et al. 2013, *ApJ*, 762, 83
 Cimatti, A., Cassata, P., Pozzetti, L., et al. 2008, *A&A*, 482, 21
 Conroy, C., & Wechsler, R. H. 2009, *ApJ*, 696, 620
 Daddi, E. et al. 2005, *ApJ*, 626, 680
 Damjanov, I., Abraham, R. G., Glazebrook, K., et al. 2011, *ApJ*, 739, L44
 Dekel, A., Sari, R., & Ceverino, D. 2009, *ApJ*, 703, 785
 Dekel, A., & Burkert, A. 2014, *MNRAS*, 438, 1870
 Di Matteo, T., Springel, V., & Hernquist, L. 2005, *Nature*, 433, 604
 Dullo, B. T., & Graham, A. W. 2013, *ApJ*, 768, 36
 Dutton, A. A. et al. 2011, *MNRAS*, 410, 1660
 Erb, D. K. et al. 2006, *ApJ*, 646, 107
 Faber, S. M. et al. 2007, *ApJ*, 665, 265
 Fan, L. et al. 2010, *ApJ*, 718, 1460
 Ferguson, H. C. et al. 2004, *ApJ*, 600, L107
 Franx, M., van Dokkum, P. G., Förster Schreiber, N. M., Wuyts, S., Labbé, I., & Toft, S. 2008, *ApJ*, 688, 770
 Fumagalli, M., Patel, S. G., Franx, M., et al. 2012, *ApJ*, 757, L22
 Gialalisco, M., Steidel, C. C., & Macchetto, F. D. 1996, *ApJ*, 470, 189
 Gialalisco, M., et al. 2004, *ApJ*, 600, L93
 González, V. et al. 2010, *ApJ*, 713, 115
 Grogin, N. A. et al. 2011, *ApJS*, 197, 35
 Guo, Y. et al. 2009, *MNRAS*, 398, 1129
 Guo, Y. et al. 2011, *ApJ*, 735, 18
 Holden, B. P., van der Wel, A., Rix, H.-W., & Franx, M. 2012, *ApJ*, 749, 96
 Hopkins, P. F., Hernquist, L., Cox, T. J., Keres, D., & Wuyts, S. 2009, *ApJ*, 691, 1424
 Huang, K.-H., Ferguson, H. C., Ravindranath, S., & Su, J. 2013, *ApJ*, 765, 68
 Huertas-Company, M., Mei, S., Shankar, F., et al. 2013, *MNRAS*, 428, 1715
 Ilbert, O., et al. 2010, *ApJ*, 709, 644
 Kelvin, L. S., Driver, S. P., Robotham, A. S. G., et al. 2012, *MNRAS*, 421, 1007
 Koekemoer, A. M. et al. 2011, *ApJS*, 197, 36
 Kormendy, J. 1977, *ApJ*, 218, 333
 Kormendy, J., & Kennicutt, R. C., Jr. 2004, *ARA&A*, 42, 603
 Kravtsov, A. V. 2013, *ApJ*, 764, L31
 Kriek, M. et al. 2009a, *ApJ*, 700, 221
 Kriek, M., van Dokkum, P. G., Franx, M., Illingworth, G. D., & Magee, D. K. 2009b, *ApJ*, 705, L71
 Labbé, I. et al. 2006, *ApJ*, 649, L67
 Laidler, V. G., Papovich, C., Grogin, N. A., et al. 2007, *PASP*, 119, 1325
 Lilly, S. et al. 1998, *ApJ*, 500, 75
 Macciò, A. V., Dutton, A. A., & van den Bosch, F. C. 2008, *MNRAS*, 391, 1940
 Maller, A. H., & Dekel, A. 2002, *MNRAS*, 335, 487
 Mancini, C. et al. 2010, *MNRAS*, 401, 933
 McGrath, E. J., Stockton, A., Canalizo, G., Iye, M., & Maihara, T. 2008, *ApJ*, 682, 303
 Mo, H. J., Mao, S., & White, S. D. M. 1998, *MNRAS*, 295, 319
 Mosleh, M. et al. 2012, *ApJ*, 756, L12
 Moster, B. P. et al. 2010, *ApJ*, 710, 903
 Moster, B. P., Naab, T., & White, S. D. M. 2013, *MNRAS*, 428, 3121
 Muzzin, A., Marchesini, D., Stefanon, M., et al. 2013, *ApJ*, 777, 18
 Naab, T., Johansson, P. H., & Ostriker, J. P. 2009, *ApJ*, 699, L178
 Newman, A. B., Ellis, R. S., Treu, T., & Bundy, K. 2010, *ApJ*, 717, L103

- Newman, A. B., Ellis, R. S., Bundy, K., & Treu, T. 2012, *ApJ*, 746, 162
- Nipoti, C., Treu, T., Leauthaud, A., et al. 2012, *MNRAS*, 422, 1714
- Oesch, P. A. et al. 2010, *ApJ*, 709, L21
- Oser, L., Naab, T., Ostriker, J. P., & Johansson, P. H. 2012, *ApJ*, 744, 63
- Pacifici, C., Kassim, S. A., Weiner, B., Charlot, S., & Gardner, J. P. 2013, *ApJ*, 762, L15
- Papovich, C., Finkelstein, S. L., Ferguson, H. C., Lotz, J. M., & Giavalisco, M. 2011, *MNRAS*, 412, 1123
- Peebles, P. J. E. 1980, *The Large-Scale Structure of the Universe* (Princeton: Princeton Univ. Press)
- Peng, Y.-j. et al. 2010, *ApJ*, 721, 193
- Poggianti, B. M. et al. 2013, *ApJ*, 762, 77
- Quadri, R. F., & Williams, R. J. 2010, *ApJ*, 725, 794
- Ravindranath, S. et al. 2004, *ApJ*, 604, L9
- Saracco, P., Longhetti, M., & Gargiulo, A. 2011, *MNRAS*, 412, 2707
- Shen, S. et al. 2003, *MNRAS*, 343, 978
- Skelton, R. E. et al. 2014, *arXiv:1403.3689*
- Somerville, R. S. et al. 2008, *ApJ*, 672, 776
- Sonnenfeld, A., Treu, T., Gavazzi, R., et al. 2013, *ApJ*, 777, 98
- Stockton, A., McGrath, E., Canalizo, G., Iye, M., & Maihara, T. 2008, *ApJ*, 672, 146
- Szomoru, D. et al. 2010, *ApJ*, 714, L244
- Szomoru, D. et al. 2011, *ApJ*, 735, L22
- Szomoru, D., Franx, M., & van Dokkum, P. G. 2012, *ApJ*, 749, 121
- Taylor, E. N. et al. 2009, *ApJS*, 183, 295
- Taylor, E. N. et al. 2010, *ApJ*, 720, 723
- Toft, S. et al. 2007, *ApJ*, 671, 285
- Toft, S. et al. 2012, *ApJ*, 754, 3
- Toft, S. et al. 2014, *ApJ*, 782, 68
- Toomre, A., & Toomre, J. 1972, *ApJ*, 178, 623
- Trager, S. C., Faber, S. M., Worthey, G., & González, J. J. 2000, *AJ*, 120, 165
- Treu, T. et al. 2005, *ApJ*, 633, 174
- Trujillo, I. et al. 2006a, *ApJ*, 650, 18
- Trujillo, I. et al. 2006b, *MNRAS*, 373, L36
- Trujillo, I., Conselice, C. J., Bundy, K., et al. 2007, *MNRAS*, 382, 109
- Trujillo, I. et al. 2009, *ApJ*, 692, L118
- Valentinuzzi, T. et al. 2010, *ApJ*, 712, 226
- van den Bosch, F. C. et al. 2008, *MNRAS*, 387, 79
- van der Wel, A. et al. 2006, *ApJ*, 652, 97
- van der Wel, A., Holden, B. P., Zirm, A. W., Franx, M., Rettura, A., Illingworth, G. D., & Ford, H. C. 2008, *ApJ*, 688, 48
- van der Wel, A., Bell, E. F., Holden, B. P., Skibba, R. A., & Rix, H.-W. 2010, *ApJ*, 714, 1779
- van der Wel, A. et al. 2012, *ApJS*, 203, 24
- van der Wel, A., Bell, E. F., van den Bosch, F. C., Gallazzi, A., & Rix, H.-W. 2009a, *ApJ*, 698, 1232
- van den Bosch, R. C. E. et al. 2012, *Nature*, 491, 729
- van der Wel, A., Rix, H.-W., Holden, B. P., Bell, E. F., & Robaina, A. R. 2009b, *ApJ*, 706, L120
- van der Wel, A. et al. 2011a, *ApJ*, 730, 38
- van der Wel, A., Straughn, A. N., Rix, H.-W., et al. 2011b, *ApJ*, 742, 111
- van der Wel, A. et al. 2013, *ApJ*, 777, L17
- van de Sande, J. et al. 2011, *ApJ*, 736, L9
- van de Sande, J. et al. 2013, *ApJ*, 771, 85
- van Dokkum, P. G. et al. 2008, *ApJ*, 677, L5
- van Dokkum, P. G. et al. 2010, *ApJ*, 709, 1018
- Wake, D. A. et al. 2011, *ApJ*, 728, 46
- Whitaker, K. E. et al. 2011, *ApJ*, 735, 86
- Williams, R. J., Quadri, R. F., Franx, M., van Dokkum, P., & Labbé, I. 2009, *ApJ*, 691, 1879
- Williams, R. J. et al. 2010, *ApJ*, 713, 738
- Williams, C. C., Giavalisco, M., Cassata, P., et al. 2014, *ApJ*, 780, 1
- Wuyts, S. et al. 2007, *ApJ*, 655, 51
- Wuyts, S. et al. 2010, *ApJ*, 722, 1666
- Wuyts, S. et al. 2012, *ApJ*, 753, 114
- Zirm, A. W. et al. 2007, *ApJ*, 656, 66

APPENDIX

A. COMPLEMENTARY SIZE DISTRIBUTIONS

Throughout this paper we use the radius R_{eff} as measured along the major axis. Circularized sizes ($R_{\text{eff,circ}} = R_{\text{eff}} \sqrt{b/a}$, where b/a is the projected axis ratio) have often been used in the literature. For this reason we provide the circularized size distributions for the early- and late-type samples in Table A1. We stress that since galaxies are predominantly oblate, that is, disk-like, using R_{eff} instead of $R_{\text{eff,circ}}$ is more prudent: R_{eff} is (almost) independent of inclination, while $R_{\text{eff,circ}}$ depends on the short, projected axis, which obviously strongly varies with inclination.

Throughout the paper we distinguish between late- and early-type galaxies on the basis of star formation activity. For some purposes it may be more useful to work with the size distributions of the full sample, without separating by type. In Table A3 we provide the size distributions of the full sample.

Finally, since stellar mass is a derived, model-dependent quantity that is potentially suffering from large systematic errors, one might be interested in galaxy size as a function of luminosity, which is essentially a directly observed quantity. In Table A3 and A2 we provide the size distributions as a function of rest-frame V -band luminosity.

Table A1

Logarithmic size distribution (16% – 84% range) as a function of galaxy mass and redshift. Identical to Table 2, but with circularized sizes instead of semi-major axis sizes.

z		Early-Type Galaxies					Late-Type Galaxies				
		$M_* = 10^{9.25}$	9.75	10.25	10.75	11.25	9.25	9.75	10.25	10.75	11.25
0.25	16%	-0.07±0.08	-0.05±0.04	0.02±0.03	0.29±0.04	...	0.07±0.01	0.21±0.02	0.27±0.02	0.51±0.03	...
	50%	0.17±0.01	0.17±0.03	0.24±0.04	0.57±0.03	0.72±0.07	0.30±0.01	0.42±0.01	0.49±0.03	0.70±0.02	0.91±0.04
	84%	0.31±0.03	0.39±0.05	0.48±0.04	0.84±0.05	1.01±0.11	0.54±0.01	0.62±0.02	0.73±0.02	0.86±0.05	1.06±0.06
0.75	16%	-0.11±0.04	-0.25±0.02	-0.11±0.02	0.11±0.02	0.57±0.02	0.04±0.01	0.17±0.01	0.27±0.01	0.39±0.01	0.64±0.04
	50%	0.12±0.02	0.10±0.02	0.13±0.02	0.36±0.01	0.74±0.03	0.26±0.00	0.39±0.01	0.49±0.01	0.61±0.02	0.82±0.03
	84%	0.34±0.02	0.34±0.03	0.33±0.01	0.56±0.01	0.93±0.04	0.47±0.01	0.58±0.01	0.66±0.01	0.78±0.01	0.98±0.02
1.25	16%	...	-0.21±0.02	-0.23±0.01	-0.07±0.02	0.32±0.04	-0.05±0.01	0.10±0.01	0.23±0.01	0.36±0.01	0.51±0.01
	50%	...	0.06±0.03	-0.01±0.02	0.16±0.02	0.49±0.03	0.19±0.00	0.32±0.00	0.42±0.00	0.53±0.01	0.67±0.02
	84%	...	0.33±0.03	0.27±0.03	0.44±0.03	0.75±0.03	0.41±0.01	0.51±0.01	0.60±0.01	0.69±0.01	0.83±0.01
1.75	16%	...	-0.12±0.06	-0.36±0.02	-0.14±0.02	0.11±0.06	-0.09±0.01	0.02±0.01	0.16±0.01	0.24±0.02	0.41±0.03
	50%	...	0.12±0.04	-0.05±0.04	0.08±0.02	0.37±0.03	0.15±0.00	0.24±0.01	0.38±0.01	0.47±0.01	0.58±0.02
	84%	...	0.36±0.04	0.27±0.04	0.42±0.03	0.67±0.03	0.36±0.01	0.46±0.01	0.55±0.01	0.64±0.01	0.75±0.02
2.25	16%	-0.53±0.09	-0.34±0.03	0.02±0.05	...	-0.06±0.01	0.06±0.02	0.16±0.02	0.29±0.04
	50%	-0.12±0.05	-0.02±0.03	0.25±0.05	...	0.17±0.01	0.29±0.01	0.39±0.01	0.52±0.02
	84%	0.25±0.03	0.41±0.06	0.72±0.12	...	0.38±0.01	0.49±0.01	0.56±0.01	0.70±0.02
2.75	16%	-0.37±0.05	-0.08±0.11	0.01±0.02	0.08±0.05	0.24±0.03
	50%	-0.01±0.04	0.29±0.09	0.27±0.01	0.34±0.01	0.43±0.02
	84%	0.40±0.06	0.62±0.10	0.48±0.02	0.54±0.02	0.63±0.06

Table A2

Logarithmic size distributions (16% – 84% range) as a function of rest-frame V -band luminosity and redshift.

z		Early-Type Galaxies					Late-Type Galaxies				
		$L_* = 10^{9.25}$	9.75	10.25	10.75	11.25	9.25	9.75	10.25	10.75	11.25
0.25	16%	0.04±0.05	0.06±0.03	0.21±0.05	0.61±0.07	...	0.16±0.03	0.31±0.02	0.42±0.02	0.68±0.06	...
	50%	0.27±0.02	0.28±0.02	0.45±0.02	0.72±0.03	...	0.42±0.02	0.57±0.01	0.69±0.02	0.92±0.03	...
	84%	0.44±0.04	0.51±0.04	0.72±0.05	0.99±0.06	...	0.64±0.02	0.76±0.01	0.87±0.02	1.07±0.04	...
0.75	16%	-0.01±0.05	-0.08±0.03	0.05±0.02	0.31±0.01	0.54±0.05	0.06±0.02	0.21±0.01	0.37±0.01	0.54±0.01	0.73±0.10
	50%	0.24±0.02	0.21±0.02	0.27±0.02	0.54±0.02	0.81±0.06	0.33±0.01	0.46±0.01	0.60±0.01	0.75±0.01	0.93±0.08
	84%	0.41±0.02	0.46±0.02	0.47±0.02	0.81±0.02	1.03±0.06	0.57±0.02	0.67±0.00	0.79±0.01	0.93±0.02	1.12±0.08
1.25	16%	-0.10±0.12	-0.12±0.04	-0.14±0.02	0.07±0.02	0.29±0.03	0.03±0.02	0.11±0.01	0.25±0.01	0.43±0.01	0.48±0.05
	50%	0.19±0.04	0.23±0.03	0.10±0.01	0.32±0.01	0.56±0.03	0.32±0.01	0.37±0.01	0.49±0.01	0.64±0.01	0.75±0.02
	84%	0.45±0.11	0.51±0.03	0.40±0.02	0.58±0.03	0.83±0.04	0.66±0.04	0.60±0.01	0.70±0.00	0.82±0.01	0.96±0.03
1.75	16%	...	-0.04±0.08	-0.19±0.03	-0.11±0.03	0.20±0.03	0.12±0.03	0.05±0.01	0.15±0.01	0.32±0.01	0.37±0.08
	50%	...	0.27±0.03	0.09±0.02	0.15±0.02	0.37±0.02	0.43±0.04	0.32±0.01	0.40±0.01	0.56±0.01	0.70±0.03
	84%	...	0.53±0.05	0.39±0.03	0.50±0.04	0.71±0.04	0.82±0.05	0.58±0.01	0.62±0.01	0.75±0.01	0.87±0.01
2.25	16%	-0.26±0.11	-0.26±0.03	0.00±0.05	...	0.01±0.04	0.08±0.01	0.20±0.01	0.26±0.06
	50%	0.08±0.04	0.07±0.04	0.24±0.03	...	0.30±0.03	0.34±0.01	0.44±0.01	0.59±0.02
	84%	0.35±0.08	0.50±0.05	0.49±0.05	...	0.68±0.06	0.56±0.01	0.65±0.01	0.79±0.03
2.75	16%	-0.27±0.03	0.01±0.05	0.09±0.01	0.18±0.01	0.26±0.07
	50%	0.03±0.05	0.34±0.09	0.33±0.02	0.43±0.01	0.54±0.02
	84%	0.42±0.07	0.73±0.08	0.62±0.04	0.63±0.01	0.75±0.02

Table A3

Logarithmic size distributions (16% – 84% range) for the full population (early- and late-type galaxies combined) as a function of mass and redshift, and rest-frame V -band luminosity and redshift.

z		Early+Late Type Galaxies									
		$M_* = 10^{9.25}$	9.75	10.25	10.75	11.25	$L_* = 10^{9.25}$	9.75	10.25	10.75	11.25
0.25	16%	0.21±0.01	0.28±0.02	0.27±0.03	0.49±0.04	...	0.14±0.02	0.26±0.02	0.37±0.01	0.62±0.04	...
	50%	0.46±0.01	0.54±0.02	0.54±0.03	0.75±0.03	...	0.37±0.02	0.53±0.01	0.62±0.02	0.79±0.03	...
	84%	0.69±0.01	0.77±0.02	0.82±0.02	0.99±0.02	...	0.62±0.02	0.74±0.01	0.85±0.01	1.05±0.03	...
0.75	16%	0.16±0.01	0.25±0.01	0.18±0.01	0.35±0.02	0.66±0.02	0.05±0.02	0.18±0.01	0.28±0.01	0.40±0.02	0.62±0.06
	50%	0.41±0.01	0.52±0.01	0.52±0.01	0.59±0.01	0.85±0.01	0.29±0.01	0.44±0.01	0.54±0.01	0.68±0.01	0.83±0.04
	84%	0.64±0.01	0.74±0.01	0.78±0.01	0.84±0.01	1.04±0.02	0.54±0.02	0.66±0.00	0.76±0.01	0.90±0.01	1.04±0.04
1.25	16%	...	0.21±0.01	0.15±0.02	0.23±0.02	0.49±0.03	0.02±0.02	0.10±0.01	0.19±0.01	0.29±0.01	0.36±0.04
	50%	...	0.47±0.01	0.52±0.01	0.57±0.01	0.74±0.02	0.30±0.02	0.36±0.01	0.47±0.01	0.58±0.01	0.67±0.04
	84%	...	0.68±0.01	0.74±0.01	0.80±0.01	0.94±0.02	0.64±0.03	0.60±0.01	0.69±0.01	0.79±0.01	0.91±0.03
1.75	16%	...	0.15±0.01	0.12±0.02	0.09±0.01	0.34±0.02	...	0.05±0.01	0.12±0.01	0.16±0.01	0.22±0.02
	50%	...	0.42±0.01	0.48±0.01	0.48±0.01	0.64±0.03	...	0.32±0.01	0.39±0.01	0.51±0.01	0.51±0.04
	84%	...	0.65±0.01	0.69±0.01	0.74±0.01	0.83±0.02	...	0.58±0.01	0.61±0.00	0.73±0.01	0.83±0.02
2.25	16%	0.10±0.02	0.03±0.03	0.28±0.03	0.07±0.01	0.15±0.01	0.10±0.03
	50%	0.41±0.01	0.45±0.02	0.59±0.03	0.33±0.01	0.43±0.01	0.45±0.03
	84%	0.63±0.01	0.68±0.01	0.83±0.02	0.55±0.01	0.64±0.01	0.74±0.03
2.75	16%	0.01±0.03	0.27±0.06	0.11±0.02	0.12±0.06
	50%	0.43±0.01	0.52±0.03	0.41±0.01	0.52±0.03
	84%	0.65±0.02	0.75±0.04	0.62±0.01	0.74±0.02